

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/00661 A1

(51) International Patent Classification⁷: **C07D 487/04**,
A61K 31/505, A61P 17/06, 19/02, 37/06

(74) Agents: **LUMB, J., Trevor** et al.; Pfizer Inc., 235 East
42nd Street, New York, NY 10017 (US).

(21) International Application Number: PCT/IB01/00975

(22) International Filing Date: 5 June 2001 (05.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/214,287 26 June 2000 (26.06.2000) US

(71) Applicant (*for all designated States except US*): **PFIZER
PRODUCTS INC.** [US/US]; Eastern Point Road, Groton,
CT 06340 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


- *with international search report*
- *before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BLUMENKOPF,
Todd, Andrew** [US/US]; Pfizer Global Research & De-
velopment, Eastern Point Road, Groton, CT 06340 (US).
FLANAGAN, Mark, Edward [US/US]; Pfizer Global
Research & Development, Eastern Point Road, Groton,
CT 06340 (US). **MUNCHHOFF, Michael, John** [US/US];
Pfizer Global Research & Development, Eastern Point
Road, Groton, CT 06340 (US).

(54) Title: PYRROLO[2,3-d]PYRIMIDINE COMPOUNDS AS IMMUNOSUPPRESSIVE AGENTS



(57) Abstract: A compound of Formula (I) wherein R¹, R² and R³ are as defined above,
useful as inhibitors of protein kinases, such as the enzyme Janus Kinase 3.

WO 02/00661 A1

PYRROLO (2,3-D) PYRIMIDINE COMPOUNDS AS IMMUNOSUPPRESSIVE AGENTS

Background of the Invention


The present invention relates to pyrrolo[2,3-d]pyrimidine compounds which are inhibitors of protein kinases, such as the enzyme Janus Kinase 3 (hereinafter also referred to as JAK3) and as such are useful therapy as immunosuppressive agents for organ transplants, xeno transplantation, lupus, multiple sclerosis, rheumatoid arthritis, psoriasis, Type I diabetes and complications from diabetes, cancer, asthma, atopic dermatitis, autoimmune thyroid disorders, ulcerative colitis, Crohn's disease, Alzheimer's disease, Leukemia and other indications where immunosuppression would be desirable.

This invention also relates to a method of using such compounds in the treatment of the above indications in mammals, especially humans, and the pharmaceutical compositions useful therefor.

JAK3 is a member of the Janus family of protein kinases. Although the other members of this family are expressed by essentially all tissues, JAK3 expression is limited to hematopoietic cells. This is consistent with its essential role in signaling through the receptors for IL-2, IL-4, IL-7, IL-9 and IL-15 by non-covalent association of JAK3 with the gamma chain common to these multichain receptors. XSCID patient populations have been identified with severely reduced levels of JAK3 protein or with genetic defects to the common gamma chain, suggesting that immunosuppression should result from blocking signaling through the JAK3 pathway. Animal studies have suggested that JAK3 not only plays a critical role in B and T lymphocyte maturation, but that JAK3 is constitutively required to maintain T cell function. Modulation of immune activity through this novel mechanism can prove useful in the treatment of T cell proliferative disorders such as transplant rejection and autoimmune diseases.


Summary of the Invention

The present invention relates to a compound of the formula



or the pharmaceutically acceptable salt thereof; wherein


R^1 is a group of the formula



wherein y is 0, 1 or 2;

- 5 R^4 is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₁-C₆)alkylsulfonyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl wherein the alkyl, alkenyl and alkynyl groups are optionally substituted by deuterium, hydroxy, amino, trifluoromethyl, (C₁-C₄)alkoxy, (C₁-C₆)acyloxy, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, cyano, nitro, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl or (C₁-C₆)acylamino; or R^4 is (C₃-C₁₀)cycloalkyl
10 wherein the cycloalkyl group is optionally substituted by deuterium, hydroxy, amino, trifluoromethyl, (C₁-C₆)acyloxy, (C₁-C₆)acylamino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, cyano, cyano(C₁-C₆)alkyl, trifluoromethyl(C₁-C₆)alkyl, nitro, nitro(C₁-C₆)alkyl or (C₁-C₆)acylamino;

- R^5 is (C₂-C₉)heterocycloalkyl wherein the heterocycloalkyl groups must be
15 substituted by one to five groups consisting of carboxy, cyano, amino, deuterium, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo, (C₁-C₆)acyl, (C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH, (C₁-C₆)alkylamino-CO-, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl, (C₁-C₆)acyloxy(C₁-C₆)alkyl, nitro, cyano(C₁-C₆)alkyl, halo(C₁-
20 C₆)alkyl, nitro(C₁-C₆)alkyl, trifluoromethyl, trifluoromethyl(C₁-C₆)alkyl, (C₁-C₆)acylamino, (C₁-C₆)acylamino(C₁-C₆)alkyl, (C₁-C₆)alkoxy(C₁-C₆)acylamino, amino(C₁-C₆)acyl, amino(C₁-C₆)acyl(C₁-C₆)alkyl, (C₁-C₆)alkylamino(C₁-C₆)acyl, ((C₁-C₆)alkyl)₂amino(C₁-C₆)acyl, $R^{15}R^{16}N-CO-O-$, $R^{15}R^{16}N-CO-(C_1-C_6)alkyl$, (C₁-C₆)alkyl-S(O)_m, $R^{15}R^{16}NS(O)_m$, $R^{15}R^{16}NS(O)_m(C_1-C_6)alkyl$, $R^{15}S(O)_mR^{16}N$, $R^{15}S(O)_mR^{16}N(C_1-$
25 C₆)alkyl wherein m is 0, 1 or 2 and R^{15} and R^{16} are each independently selected from hydrogen or (C₁-C₆)alkyl; and a group of the formula



II

wherein a is 0, 1, 2, 3 or 4;

b, c, e, f and g are each independently 0 or 1;

d is 0, 1, 2, or 3;

5 X is $S(O)_n$ wherein n is 0, 1 or 2; oxygen, carbonyl or $-C(=N\text{-cyano})-$;

Y is $S(O)_n$ wherein n is 0, 1 or 2; or carbonyl; and

Z is carbonyl, $C(O)O-$, $C(O)NR-$ wherein R is hydrogen or (C_1-C_6) alkyl; or Z is $S(O)_n$ wherein n is 0, 1 or 2;

10 R^6 , R^7 , R^8 , R^9 , R^{10} and R^{11} are each independently selected from the group consisting of hydrogen or (C_1-C_6) alkyl optionally substituted by deuterium, hydroxy, amino, trifluoromethyl, (C_1-C_6) acyloxy, (C_1-C_6) acylamino, (C_1-C_6) alkylamino, $((C_1-C_6)alkyl)_2$ amino, cyano, cyano (C_1-C_6) alkyl, trifluoromethyl (C_1-C_6) alkyl, nitro, nitro (C_1-C_6) alkyl or (C_1-C_6) acylamino;

15 R^{12} is (C_6-C_{10}) aryl, (C_2-C_9) heteroaryl, (C_3-C_{10}) cycloalkyl or (C_2-C_9) heterocycloalkyl, wherein the aryl, heteroaryl, cycloalkyl and heterocycloalkyl groups are optionally substituted by one to four groups consisting of hydrogen, deuterium, amino, halo, oxo, hydroxy, nitro, carboxy, (C_2-C_6) alkenyl, (C_2-C_6) alkynyl, trifluoromethyl, trifluoromethoxy, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, (C_3-C_{10}) cycloalkyl, $(C_1-C_6)alkyl-CO-NH-$, $(C_1-C_6)alkoxy-CO-NH-$, $(C_1-C_6)alkyl-CO-NH-(C_1-C_6)alkyl$, $(C_1-C_6)alkoxy-CO-NH-(C_1-C_6)alkyl$, $(C_1-C_6)alkoxy-CO-NH-(C_1-C_6)alkoxy$, carboxy, carboxy $(C_1-C_6)alkyl$, carboxy $(C_1-C_6)alkoxy$, benzyloxycarbonyl $(C_1-C_6)alkoxy$, $(C_1-C_6)alkoxycarbonyl(C_1-C_6)alkoxy$, (C_6-C_{10}) aryl, amino, amino $(C_1-C_6)alkyl$, $(C_1-C_6)alkoxycarbonylamino$, $(C_6-C_{10})aryl(C_1-C_6)alkoxycarbonylamino$, $(C_1-C_6)alkylamino$, $((C_1-C_6)alkyl)_2$ amino, $(C_1-C_6)alkylamino(C_1-C_6)alkyl$, $((C_1-C_6)alkyl)_2$ amino $(C_1-C_6)alkyl$,
 20 hydroxy, $(C_1-C_6)alkoxy$, carboxy, carboxy $(C_1-C_6)alkyl$, $(C_1-C_6)alkoxycarbonyl$, $(C_1-C_6)alkoxycarbonyl(C_1-C_6)alkyl$, $(C_1-C_6)alkoxy-CO-NH-$, $(C_1-C_6)alkyl-CO-NH-$, cyano, (C_5-C_9) heterocycloalkyl, amino- $CO-NH-$, $(C_1-C_6)alkylamino-CO-NH-$, $((C_1-C_6)alkyl)_2$ amino- $CO-NH-$, $(C_6-C_{10})aryl$ amino- $CO-NH-$, (C_5-C_9) heteroarylamino- $CO-$
 25

- NH-, (C₁-C₆)alkylamino-CO-NH-(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino-CO-NH-(C₁-C₆)alkyl, (C₆-C₁₀)arylamino-CO-NH-(C₁-C₆)alkyl, (C₅-C₉)heteroarylamino-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkylcyano, (C₁-C₆)alkylcarboxy(C₁-C₆)alkoxy, (C₁-C₆)alkylcarboxy, sulfonylamino, aminosulfonyl, sulfonylamino(C₁-C₆)alkyl, sulfonylaminocarboxy(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₆-C₁₀)arylsulfonyl, (C₆-C₁₀)arylsulfonylamino, (C₆-C₁₀)arylsulfonylamino(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₃-C₁₀)cycloalkyl, (C₃-C₁₀)cycloalkoxy, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₆-C₁₀)arylamino, (C₁-C₆)alkylthio, (C₆-C₁₀)arylthio, (C₁-C₆)alkylsulfinyl, (C₆-C₁₀)arylsulfinyl, (C₁-C₆)alkylsulfonyl, (C₆-C₁₀)arylsulfonyl, (C₁-C₆)acyl, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkylamino-CO-, (C₅-C₉)heteroaryl, (C₂-C₉)heterocycloalkyl or (C₆-C₁₀)aryl wherein the heteroaryl, heterocycloalkyl and aryl groups which are optionally substituted on R¹² may be further substituted by one to three groups consisting of halo, (C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-NH-, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkoxy, carboxy, carboxy(C₁-C₆)alkyl, carboxy(C₁-C₆)alkoxy, benzyloxycarbonyl(C₁-C₆)alkoxy, (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkoxy, (C₆-C₁₀)aryl, amino, amino(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonylamino, (C₆-C₁₀)aryl(C₁-C₆)alkoxycarbonylamino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₁-C₆)alkylamino(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino(C₁-C₆)alkyl, hydroxy, (C₁-C₆)alkoxy, carboxy, carboxy(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-, cyano, (C₅-C₉)heterocycloalkyl, amino-CO-NH-, (C₁-C₆)alkylamino-CO-NH-, ((C₁-C₆)alkyl)₂amino-CO-NH-, (C₆-C₁₀)arylamino-CO-NH-, (C₅-C₉)heteroarylamino-CO-NH-, (C₁-C₆)alkylamino-CO-NH-(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino-CO-NH-(C₁-C₆)alkyl, (C₆-C₁₀)arylamino-CO-NH-(C₁-C₆)alkyl, (C₅-C₉)heteroarylamino-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₆-C₁₀)arylsulfonyl, (C₆-C₁₀)arylsulfonylamino, (C₆-C₁₀)arylsulfonylamino(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₅-C₉)heteroaryl and (C₂-C₉)heterocycloalkyl;
- R² and R³ are each independently selected from the group consisting of hydrogen, deuterium, amino, halo, hydroxy, nitro, carboxy, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, trifluoromethyl, trifluoromethoxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₃-C₁₀)cycloalkyl wherein the alkyl, alkoxy or cycloalkyl groups are optionally substituted by one to three groups selected from halo, hydroxy, carboxy, amino (C₁-C₆)alkylthio,

(C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₅-C₉)heteroaryl, (C₂-C₉)heterocycloalkyl, (C₃-C₉)cycloalkyl or (C₆-C₁₀)aryl; or R² and R³ are each independently (C₃-C₁₀)cycloalkyl, (C₃-C₁₀)cycloalkoxy, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₆-C₁₀)arylamino, (C₁-C₆)alkylthio, (C₆-C₁₀)arylthio, (C₁-C₆)alkylsulfinyl, (C₆-C₁₀)arylsulfinyl, (C₁-C₆)alkylsulfonyl, (C₆-C₁₀)arylsulfonyl, (C₁-C₆)acyl, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkylamino-CO-, (C₅-C₉)heteroaryl, (C₂-C₉)heterocycloalkyl or (C₆-C₁₀)aryl wherein the heteroaryl, heterocycloalkyl and aryl groups are optionally substituted by one to three halo, (C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-NH-, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkoxy, carboxy, carboxy(C₁-C₆)alkyl, carboxy(C₁-C₆)alkoxy, benzyloxycarbonyl(C₁-C₆)alkoxy, (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkoxy, (C₆-C₁₀)aryl, amino, amino(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonylamino, (C₆-C₁₀)aryl(C₁-C₆)alkoxycarbonylamino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₁-C₆)alkylamino(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino(C₁-C₆)alkyl, hydroxy, (C₁-C₆)alkoxy, carboxy, carboxy(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-, cyano, (C₅-C₉)heterocycloalkyl, amino-CO-NH-, (C₁-C₆)alkylamino-CO-NH-, ((C₁-C₆)alkyl)₂amino-CO-NH-, (C₆-C₁₀)arylamino-CO-NH-, (C₅-C₉)heteroarylamino-CO-NH-, (C₁-C₆)alkylamino-CO-NH-(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino-CO-NH-(C₁-C₆)alkyl, (C₆-C₁₀)arylamino-CO-NH-(C₁-C₆)alkyl, (C₅-C₉)heteroarylamino-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₆-C₁₀)arylsulfonyl, (C₆-C₁₀)arylsulfonylamino, (C₆-C₁₀)arylsulfonylamino(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₅-C₉)heteroaryl or (C₂-C₉)heterocycloalkyl;

with the proviso that R⁵ must be substituted by the group of formula II.

The present invention also relates to the pharmaceutically acceptable acid addition salts of compounds of the formula I. The acids which are used to prepare the pharmaceutically acceptable acid addition salts of the aforementioned base compounds of this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, acetate, lactate, citrate, acid citrate, tartrate, bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate,

benzenesulfonate, p-toluenesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)]salts.

The invention also relates to base addition salts of formula I. The chemical bases that may be used as reagents to prepare pharmaceutically acceptable base salts of those compounds of formula I that are acidic in nature are those that form non-toxic base salts with such compounds. Such non-toxic base salts include, but are not limited to those derived from such pharmacologically acceptable cations such as alkali metal cations (e.g., potassium and sodium) and alkaline earth metal cations (e.g., calcium and magnesium), ammonium or water-soluble amine addition salts such as N-methylglucamine-(meglumine), and the lower alkanolammonium and other base salts of pharmaceutically acceptable organic amines.

The term "Oxone®" is a name of a monopersulfate compound used in this invention, having the formula $2\text{KHSO}_5 \cdot \text{KHSO}_4 \cdot \text{K}_2\text{SO}_4$, and sold by Aldrich Chemical Company, P.O. Box 2060, Milwaukee, WI 53201, USA.

The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight or branched moieties or combinations thereof.

The term "alkoxy", as used herein, includes O-alkyl groups wherein "alkyl" is defined above.

The term "halo", as used herein, unless otherwise indicated, includes fluoro, chloro, bromo or iodo.

The compounds of this invention may contain double bonds. When such bonds are present, the compounds of the invention exist as cis and trans configurations and as mixtures thereof.

Unless otherwise indicated, the alkyl and alkenyl groups referred to herein, as well as the alkyl moieties of other groups referred to herein (e.g., alkoxy), may be linear or branched, and they may also be cyclic (e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl) or be linear or branched and contain cyclic moieties. Unless otherwise indicated, halogen includes fluorine, chlorine, bromine, and iodine.

(C₂-C₉)Heterocycloalkyl when used herein refers to pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydropyranyl, pyranyl, thiopyranyl, aziridinyl, oxiranyl, methylenedioxy, chromenyl, isoxazolidinyl, 1,3-oxazolidin-3-yl, isothiazolidinyl, 1,3-thiazolidin-3-yl, 1,2-pyrazolidin-2-yl, 1,3-pyrazolidin-1-yl,

piperidinyl, thiomorpholinyl, 1,2-tetrahydrothiazin-2-yl, 1,3-tetrahydrothiazin-3-yl, tetrahydrothiadiazinyl, morpholinyl, 1,2-tetrahydrodiazin-2-yl, 1,3-tetrahydrodiazin-1-yl, tetrahydroazepinyl, piperazinyl, chromanyl, etc. One of ordinary skill in the art will understand that the connection of said (C₂-C₉)heterocycloalkyl rings is through a
5 carbon or a sp³ hybridized nitrogen heteroatom.

(C₂-C₉)Heteroaryl when used herein refers to furyl, thienyl, thiazolyl, pyrazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrrolyl, triazolyl, tetrazolyl, imidazolyl, 1,3,5-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,3-oxadiazolyl, 1,3,5-thiadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, 1,2,4-triazinyl, 1,2,3-
10 triazinyl, 1,3,5-triazinyl, pyrazolo[3,4-b]pyridinyl, cinnolinyl, pteridinyl, purinyl, 6,7-dihydro-5H-[1]pyrindinyl, benzo[b]thiophenyl, 5, 6, 7, 8-tetrahydro-quinolin-3-yl, benzoxazolyl, benzothiazolyl, benzisothiazolyl, benzisoxazolyl, benzimidazolyl, thianaphthenyl, isothianaphthenyl, benzofuranyl, isobenzofuranyl, isoindolyl, indolyl, indoliziny, indazolyl, isoquinolyl, quinolyl, phthalazinyl, quinoxaliny, quinazoliny,
15 benzoxazinyl; etc. One of ordinary skill in the art will understand that the connection of said (C₂-C₉)heterocycloalkyl rings is through a carbon atom or a sp³ hybridized nitrogen heteroatom.

(C₆-C₁₀)aryl when used herein refers to phenyl or naphthyl.

Compounds of formula (I) may be administered in a pharmaceutically
20 acceptable form either alone or in combination with one or more additional agents which modulate a mammalian immune system or with antiinflammatory agents. These agents may include but are not limited to cyclosporin A (e.g. Sandimmune® or Neoral®, rapamycin, FK-506 (tacrolimus), leflunomide, deoxyspergualin, mycophenolate (e.g. Cellcept®), azathioprine (e.g. Imuran®), daclizumab (e.g.
25 Zenapax®. OKT3 (e.g. Orthoclone®), AtGam, aspirin, acetaminophen, ibuprofen, naproxen, piroxicam, and antiinflammatory steroids (e.g. prednisolone or dexamethasone). These agents may be administered as part of the same or separate dosage forms, via the same or different routes of administration, and on the same or different administration schedules according to standard pharmaceutical
30 practice.

The compounds of this invention include all conformational isomers (e.g., cis and trans isomers. The compounds of the present invention have asymmetric centers and therefore exist in different enantiomeric and diastereomeric forms. This invention relates to the use of all optical isomers and stereoisomers of the

compounds of the present invention, and mixtures thereof, and to all pharmaceutical compositions and methods of treatment that may employ or contain them. In this regard, the invention includes both the E and Z configurations. The compounds of formula I may also exist as tautomers. This invention relates to the use of all such
5 tautomers and mixtures thereof.

This invention also encompasses pharmaceutical compositions containing prodrugs of compounds of the formula I. This invention also encompasses methods of treating or preventing disorders that can be treated or prevented by the inhibition of protein kinases, such as the enzyme Janus Kinase 3 comprising administering
10 prodrugs of compounds of the formula I. Compounds of formula I having free amino, amido, hydroxy or carboxylic groups can be converted into prodrugs. Prodrugs include compounds wherein an amino acid residue, or a polypeptide chain of two or more (e.g., two, three or four) amino acid residues which are covalently joined through peptide bonds to free amino, hydroxy or carboxylic acid groups of
15 compounds of formula I. The amino acid residues include the 20 naturally occurring amino acids commonly designated by three letter symbols and also include, 4-hydroxyproline, hydroxylysine, demosine, isodemosine, 3-methylhistidine, norvlin, beta-alanine, gamma-aminobutyric acid, citrulline, homocysteine, homoserine, ornithine and methioine sulfone. Prodrugs also include compounds wherein
20 carbonates, carbamates, amides and alkyl esters which are covalently bonded to the above substituents of formula I through the carbonyl carbon prodrug sidechain.

Preferred compounds of formula I include those wherein R^5 is (C_2-C_9) heterocycloalkyl optionally substituted by one to three groups selected from deuterium, hydroxy, (C_1-C_6) alkyl, halo, (C_1-C_6) alkoxy and a group of formula II.

25 Other preferred compounds of formula I include those wherein a is 0; b is 1; X is carbonyl; c is 0; d is 0; e is 0; f is 0; and g is 0.

Other preferred compounds of formula I include those wherein a is 0; b is 1; X is carbonyl; c is 0; d is 1; e is 0; f is 0, and g is 0.

30 Other preferred compounds of formula I include those wherein a is 0; b is 1; X is carbonyl; c is 1; d is 0; e is 0; f is 0; and g is 0.

Other preferred compounds of formula I include those wherein a is 0; b is 1; X is $-C(=N=\text{cyano})-$; c is 1; d is 0; e is 0; f is 0; and g is 0.

Other preferred compounds of formula I include those wherein a is 0; b is 0; c is 0; d is 0; e is 0; f is 0; g is 1; and Z is $-C(O)-O-$.

Other preferred compounds of formula I include those wherein a is 0; b is 1; X is $S(O)_n$; n is 2; c is 0; d is 0; e is 0; f is 0; and g is 0.

Other preferred compounds of formula I include those wherein a is 0; b is 1; X is $S(O)_n$; n is 2; c is 0; d is 2; e is 0; f is 1; g is 1; and Z is carbonyl.

5 Other preferred compounds of formula I include those wherein a is 0; b is 1; X is $S(O)_n$; n is 2; c is 0; d is 2; e is 0; f is 1; and g is 0.

Other preferred compounds of formula I include those wherein a is 0; b is 1; X is carbonyl; c is 1; d is 0; e is 1; Y is $S(O)_n$; n is 2; f is 0; and g is 0.

10 Other preferred compounds of formula I include those wherein a is 0; b is 1; X is $S(O)_n$; n is 2; c is 1; d is 0; e is 0; f is 0; and g is 0.

Other preferred compounds of formula I include those wherein a is 1; b is 1; X is carbonyl; c is 1; d is 0; e is 0; f is 0; and g is 0.

Other preferred compounds of formula I include those wherein a is 0; b is 1; X is $S(O)_n$; c is 0; d is 1; e is 1; Y is $S(O)_n$; n is 2; f is 0; and g is 0.

15 Other preferred compounds of formula I include those wherein a is 0; b is 1; X is $S(O)_n$; c is 0; d is 1; e is 1; Y is $S(O)_n$; n is 2; f is 1; and g is 0.

Other preferred compounds of formula I include those wherein a is 0; b is 1; X is oxygen; c is 0; d is 1; e is 1; Y is $S(O)_n$; n is 2; f is 1; and g is 0.

20 Other preferred compounds of formula I include those wherein a is 0; b is 1; X is oxygen; c is 0; d is 1; e is 1; Y is $S(O)_n$; n is 2; f is 0; and g is 0.

Other preferred compounds of formula I include those wherein a is 0; b is 1; X is carbonyl; c is 1; d is 1; e is 1; Y is $S(O)_n$; f is 0; and g is 0.

Other preferred compounds of formula I include those wherein a is 0; b is 1; X is carbonyl; c is 1; d is 1; e is 1; Y is $S(O)_n$; n is 2; f is 1; and g is 0.

25 Other preferred compounds of formula I include those wherein R^{12} is (C_6-C_{10}) aryl or (C_2-C_9) heteroaryl wherein the aryl or heteroaryl group is optionally substituted by one to four groups consisting of hydrogen, halo, hydroxy, carboxy, trifluormethyl, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, (C_1-C_6) alkyl-CO-NH-, amino, amino (C_1-C_6) alkyl, (C_1-C_6) alkylamino, $((C_1-C_6)alkyl)_2$ amino, cyano, amino-CO-NH-, (C_1-C_6) alkylamino-CO-NH-, $((C_1-C_6)alkyl)_2$ amino-CO-NH-, (C_5-C_9) heteroaryl-amino-CO-NH-, (C_1-C_6) alkylsulfonyl, (C_1-C_6) alkylsulfonylamino, (C_6-C_{10}) arylsulfonylamino, (C_1-C_6) alkylsulfonylamino, and (C_1-C_6) alkoxy-CO-NH-.

30

Specific preferred compounds of formula I include those wherein said compound is selected from the group consisting of:

- 4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-ylmethyl}-benzenesulfonamide;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-sulfamoyl-phenyl)-amide;
- 5 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-nitro-phenyl)-amide;
- {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-tetrazol-1-yl-ethanone;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methylsulfamoyl-phenyl)-amide;
- 10 (3-Hydroxy-pyrrolidin-1-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone;
- [2-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-thiazol-4-yl]-acetic acid;
- 15 Methyl-(4-methyl-5'-nitro-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;
- 5-(2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-oxo-ethyl)-thiazolidine-2,4-dione;
- {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-thiazolidin-3-yl-methanone;
- 20 Methyl-[4-methyl-1-(5-nitro-thiazol-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;
- [2-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-thiazol-4-yl]-acetic acid ethyl ester;
- 25 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methanesulfonyl-phenyl)-amide;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid thiazol-2-ylamide;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-cyano-phenyl)-amide;
- 30 {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-pyrrolidin-1-yl-methanone;
- Furan-2-carboxylic acid (2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide;

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}
(tetrahydro-furan-3-yl)-methanone;

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid isoxazol-3-ylamide;

5 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (6-cyano-pyridin-3-yl)-amide;

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-
tetrahydro-2H-[1,2']bipyridinyl-5'-carbonitrile

10 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (4-methyl-thiazol-2-yl)-amide;

2-Cyclopropyl-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-
amino]-piperidin-1-yl}-ethanone;

Cyclopentyl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-
piperidin-1-yl}-Methanone;

15 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (3-methyl-isoxazol-4-yl)-amide;

[4-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-
1-carbonyl}-amino)-phenyl]-acetic acid;

20 [1-(5-Amino-thiazol-2-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-
d]pyrimidin-4-yl)-amine;

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (3-methyl-isothiazol-5-yl)-amide;

3-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carbonyl}-cyclopentanone;

25 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid benzyl-methyl-amide; and

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid dimethylamide.

30 The present invention also relates to a pharmaceutical composition for (a)
treating or preventing a disorder or condition selected from organ transplant rejection,
xeno transplation, lupus, multiple sclerosis, rheumatoid arthritis, psoriasis, Type I
diabetes and complications from diabetes, cancer, asthma, atopic dermatitis,
autoimmune thyroid disorders, ulcerative colitis, Crohn's disease, Alzheimer's
disease, Leukemia, and other autoimmune diseases or (b) the inhibition of protein

kinases or Janus Kinase 3 (JAK3) in a mammal, including a human, comprising an amount of a compound of formula I or a pharmaceutically acceptable salt thereof, effective in such disorders or conditions and a pharmaceutically acceptable carrier.


5 The present invention also relates to a method for the inhibition of protein tyrosine kinases or Janus Kinase 3 (JAK3) in a mammal, including a human, comprising administering to said mammal an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.





10 The present invention also relates to a method for treating or preventing a disorder or condition selected from organ transplant rejection, xeno transplation, lupus, multiple sclerosis, rheumatoid arthritis, psoriasis, Type I diabetes and complications from diabetes, cancer, asthma, atopic dermatitis, autoimmune thyroid disorders, ulcerative colitis, Crohn's disease, Alzheimer's disease, Leukemia, and other autoimmune diseases in a mammal, including a human, comprising administering to said mammal an amount of a compound of formula I or a
15 pharmaceutically acceptable salt thereof, effective in treating such a condition.


Detailed Description of the Invention

The following reaction Schemes illustrate the preparation of the compounds of the present invention. Unless otherwise indicated R^2 , R^3 , R^4 and R^5 in the reaction Schemes and the discussion that follow are defined as above.



5

PREPARATION A



PREPARATION B

PREPARATION C



XXXI



XXX











XXIX



XXVIII


SCHEME 1



SCHEME 2

SCHEME 3

5



In reaction 1 of Preparation A, the 4-chloropyrrolo[2,3-d]pyrimidine compound of formula **XXI**, wherein R is hydrogen or a protecting group such as benzenesulfonyl or benzyl, is converted to the 4-chloro-5-halopyrrolo[2,3-d]pyrimidine compound of formula **XX**, wherein Y is chloro, bromo or iodo, by reacting **XXI** with N-chlorosuccinimide, N-bromosuccinimide or N-iodosuccinimide. The reaction mixture is heated to reflux, in chloroform, for a time period between about 1 hour to about 3 hours, preferably about 1 hour. Alternatively, in reaction 1 of Preparation A, the 4-chloropyrrolo[2,3-d]pyrimidine of formula **XXI**, wherein R is hydrogen, is converted to the corresponding 4-chloro-5-nitropyrrolo[2,3-d]pyrimidine of formula **XX**, wherein Y is nitro, by reacting **XXI** with nitric acid in sulfuric acid at a temperature between about -10°C to about 10°C, preferably about 0°C, for a time period between about 5 minutes to about 15 minutes, preferably about 10 minutes. The compound of formula **XXI**, wherein Y is nitro, is converted to the corresponding 4-chloro-5-aminopyrrolo[2,3-d]pyrimidine of the formula **XX**, wherein Y is amino, by reacting **XXI** under a variety of conditions known to one skilled in the art such as palladium hydrogenolysis or tin(IV)chloride and hydrochloric acid.

In reaction 2 of Preparation A, the 4-chloro-5-halopyrrolo[2,3-d]pyrimidine compound of formula **XX**, wherein R is hydrogen, is converted to the corresponding compound of formula **XIX**, wherein R² is (C₁-C₆)alkyl or benzyl, by treating **XX** with N-butyllithium, at a temperature of about -78°C, and reacting the dianion intermediate so formed with an alkylhalide or benzylhalide at a temperature between about -78°C to room temperature, preferably room temperature. Alternatively, the dianion so formed is reacted with molecular oxygen to form the corresponding 4-chloro-5-hydroxypyrrolo[2,3-d]pyrimidine compound of formula **XIX**, wherein R² is hydroxy. The compound of formula **XX**, wherein Y is bromine or iodine and R is benzenesulfonate, is converted to the compound of formula **XIX**, wherein R² is (C₆-C₁₂)aryl or vinyl, by treating **XX** with N-butyllithium, at a temperature of about -78°C, followed by the addition of zinc chloride, at a temperature of about -78°C. The corresponding organo zinc intermediate so formed is then reacted with aryl iodide or vinyl iodide in the presence of a catalytic quantity of palladium. The reaction mixture is stirred at a temperature between about 50°C to about 80°C, preferably about 70°C, for a time period between about 1 hour to about 3 hours, preferably about 1 hour.

In reaction 3 of Preparation A, the compound of formula **XIX** is converted to the corresponding compound of formula **XVI** by treating **XIX** with N-butyllithium, lithium diisopropylamine or sodium hydride, at a temperature of about -78°C, in the presence of a polar aprotic solvent, such as tetrahydrofuran. The anionic intermediate so formed is further reacted with (a) alkylhalide or benzylhalide, at a temperature between about -78°C to room temperature, preferably -78°C, when R³ is alkyl or benzyl; (b) an aldehyde or ketone, at a temperature between about -78°C to room temperature, preferably -78°C, when R³ is alkoxy; and (c) zinc chloride, at a temperature between about -78°C to room temperature, preferably -78°C, and the corresponding organozinc intermediate so formed is then reacted with aryl iodide or vinyl iodide in the presence of a catalytic quantity of palladium. The resulting reaction mixture is stirred at a temperature between about 50°C to about 80°C, preferably about 70°C, for a time period between about 1 hour to about 3 hours, preferably about 1 hour. Alternatively, the anion so formed is reacted with molecular oxygen to form the corresponding 4-chloro-6-hydroxypyrrolo[2,3-d]pyrimidine compound of formula **XVI**, wherein R³ is hydroxy.

In reaction 1 of Preparation B, the 4-chloropyrrolo[2,3-d]pyrimidine compound of formula **XXI** is converted to the corresponding compound of formula **XXII**, according to a procedure analogous to that described above in reaction 3 of Preparation A.

In reaction 2 of Preparation B, the compound of formula **XXII** is converted to the corresponding compound of formula **XVI**, according to procedures analogous to that described above in reactions 1 and 2 of Preparation A.

In reaction 1 of Preparation C, the 4-methylpyridine compound of formula **XXXI** is converted to the corresponding compound of formula **XXX** by first alkylating **XXXI** with benzylchloride in the presence of a polar aprotic solvent, such as acetone. The reaction mixture is stirred at a temperature between about 40°C to about 80°C for a time period between about 4 hours to about 24 hours. The pyridinium intermediate so formed is then reduced with a reducing agent, such as sodium borohydride, in the presence of a polar protic solvent, such as methanol, ethanol, water or mixtures thereof. The reaction is stirred at a temperature between about 0°C to a about room temperature, for a time period between about 18 hours to 24 hours.

In reaction 2 of Preparation C, the compound of formula **XXX** is converted to the corresponding compound of formula **XXIX** by treating **XXX** with borotrifluoride etherate in the presence of a reducing agent and an aprotic solvent, such as tetrahydrofuran. The reaction mixture is stirred at a temperature between about 0°C to room temperature, for a time period between about 1 hour to about 3 hours. The intermediate complex so formed is then basified with aqueous sodium hydroxide and then treated with an oxidizing agent, such as hydrogen peroxide or Oxone®, at a temperature between about 0°C to room temperature, for a time period between about 12 hours to about 24 hours.

In reaction 3 of Preparation C, the compound of formula **XXIX** is treated with an oxidizing agent, such as chromium oxide or dimethylsulfoxide, oxalylchloride or SO₃-pyridine complex, for a time period between about 1 hour to 3 hours, at ambient temperature. The ketone intermediate so formed, is then treated with an amine (R⁴-NH₂) in the presence of an acid, such as acetic acid, at about room temperature, for a time period between about 2 to about 24 hours, in an organic solvent such as methanol, ethanol or tetrahydrofuran. The corresponding imine intermediate so formed is then treated with a reducing agent, such as sodium borohydride or sodium cyanoborohydride or sodium triacetoxymethylborohydride, at ambient temperature, for a time period about 2 to about 24 hours.

In reaction 1 of Scheme 1, the 4-chloropyrrolo[2,3-d]pyrimidine compound of formula **XVII** is converted to the corresponding compound of formula **XVI**, wherein R is benzenesulfonyl or benzyl, by treating **XVII** with benzenesulfonyl chloride, benzylchloride or benzylbromide in the presence of a base, such as sodium hydride or potassium carbonate, and a polar aprotic solvent, such as dimethylformamide or tetrahydrofuran. The reaction mixture is stirred at a temperature between about 0°C to about 70°C, preferably about 30°C, for a time period between about 1 hour to about 3 hours, preferably about 2 hours.

In reaction 2 of Scheme 1, the 4-chloropyrrolo[2,3-d]pyrimidine compound of formula **XVI** is converted to the corresponding 4-aminopyrrolo[2,3-d]pyrimidine compound of formula **XV** by coupling **XVI** with an amine of the formula HNR⁴R⁵. The reaction is carried out in an alcohol solvent, such as tert-butanol, methanol or ethanol, or other high boiling organic solvents, such as dimethylformamide, triethylamine, 1,4-dioxane or 1,2-dichloroethane, at a temperature between about 60°C to about 120°C,

preferably about 80°C. Typical reaction times are between about 2 hours to about 48 hours, preferably about 16 hours. When R⁵ is a nitrogen containing heterocycloalkyl group, each nitrogen must be protected by a protecting group, such a benzyl. Removal of the R⁵ protecting group is carried out under conditions appropriate for that particular protecting group in use which will not affect the R protecting group on the pyrrolo[2,3-d]pyrimidine ring. Removal of the R⁵ protecting group, when benzyl, is carried out in an alcohol solvent, such as ethanol, in the present of hydrogen and a catalyst, such as palladium hydroxide on carbon. The R⁵ nitrogen containing heterocycloalkyl group so formed may be further reacted with a variety of different electrophiles of formula II. For urea formation, electrophiles of formula II such as isocyanates, carbamates and carbamoyl chlorides are reacted with the R⁵ nitrogen of the heteroalkyl group in a solvent, such as acetonitrile or dimethylformamide, in the presence of a base, such as sodium or potassium carbonate, at a temperature between about 20°C to about 100°C for a time period between about 24 hours to about 72 hours. For amide and sulfonamide formation, electrophiles of formula II, such as acylchlorides and sulfonyl chlorides, are reacted with the R⁵ nitrogen of the heteroalkyl group in a solvent such as methylene chloride in the presence of a base such as pyridine at ambient temperatures for a time period between about 12 hours to about 24 hours. Amide formation may also be carried out by reacting a carboxylic acid with the heteroalkyl group in the presence of a carbodiimide such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide in a solvent such as methylene chloride at ambient temperatures for 12-24 hours. For alkyl formation, electrophiles of formula II, such as α,β -unsaturated amides, acids, nitriles, esters, and α -halo amides, are reacted with the R⁵ nitrogen of the heteroalkyl group in a solvent such as methanol at ambient temperatures for a time period between about 12 hours to about 18 hours. Alkyl formation may also be carried out by reacting aldehydes with the heteroalkyl group in the presence of a reducing agent, such as sodium cyanoborohydride, in a solvent, such as methanol, at ambient temperature for a time period between about 12 hours to about 18 hours.

In reaction 3 of Scheme 1, removal of the protecting group from the compound of formula XV, wherein R is benzenesulfonyl, to give the corresponding compound of formula I, is carried out by treating XV with an alkali base, such as sodium hydroxide or potassium hydroxide, in an alcohol solvent, such as methanol or

ethanol, or mixed solvents, such as alcohol/tetrahydrofuran or alcohol/water. The reaction is carried out at room temperature for a time period between about 15 minutes to about 1 hour, preferably 30 minutes. Removal of the protecting group from the compound of formula **XV**, wherein R is benzyl, is conducted by treating **XV** with sodium in ammonia at a temperature of about -78°C for a time period between about 15 minutes to about 1 hour.

In reaction 1 of Scheme 2, the 4-chloropyrrolo[2,3-d]pyrimidine compound of formula **XX** is converted to the corresponding 4-aminopyrrolo[2,3-d]pyrimidine compound of formula **XXIV**, according to a procedure analogous to that described above in reaction 2 of Scheme 1.

In reaction 2 of Scheme 2, the 4-amino-5-halopyrrolo[2,3-d]pyrimidine compound of formula **XXIV**, wherein R is benzenesulfonate and Z is bromine or iodine, is converted to the corresponding compound of formula **XXIII** by reacting **XXIV** with (a) arylboronic acid, when R² is aryl, in an aprotic solvent, such as tetrahydrofuran or dioxane, in the presence of a catalytic quantity of palladium (0) at a temperature between about 50°C to about 100°C, preferably about 70°C, for a time period between about 2 hours to about 48 hours, preferably about 12 hours; (b) alkynes, when R² is alkynyl, in the presence of a catalytic quantity of copper (I) iodide and palladium (0), and a polar solvent, such as dimethylformamide, at room temperature, for a time period between about 1 hour to about 5 hours, preferably about 3 hours; and (c) alkenes or styrenes, when R² is vinyl or styrenyl, in the presence of a catalytic quantity of palladium in dimethylformamide, dioxane or tetrahydrofuran, at a temperature between about 80°C to about 100°C, preferably about 100°C, for a time period between about 2 hours to about 48 hours, preferably about 48 hours.

In reaction 3 of Scheme 2, the compound of formula **XXIII** is converted to the corresponding compound of formula **XV**, according to a procedure analogous to that described above in reaction 3 of Preparation A.

In reaction 1 of Scheme 3, the compound of formula **XVII** is converted to the corresponding compound of formula **I**, according to a procedure analogous to that described above in reaction 2 of Scheme 1.

The compounds of the present invention that are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids.

Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate the compound of the present invention from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained. The desired acid salt can also be precipitated from a solution of the free base in an organic solvent by adding to the solution an appropriate mineral or organic acid.

Those compounds of the present invention that are acidic in nature, are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particularly, the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the acidic compounds of the present invention. Such non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium calcium and magnesium, etc. These salts can easily be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure completeness of reaction and maximum yields of the desired final product.

The compositions of the present invention may be formulated in a conventional manner using one or more pharmaceutically acceptable carriers. Thus, the active compounds of the invention may be formulated for oral, buccal, intranasal, parenteral (e.g., intravenous, intramuscular or subcutaneous) or rectal administration

or in a form suitable for administration by inhalation or insufflation. The active compounds of the invention may also be formulated for sustained delivery.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, methyl cellulose or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters or ethyl alcohol); and preservatives (e.g., methyl or propyl p-hydroxybenzoates or sorbic acid).

For buccal administration, the composition may take the form of tablets or lozenges formulated in conventional manner.

The active compounds of the invention may be formulated for parenteral administration by injection, including using conventional catheterization techniques or infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulating agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for reconstitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The active compounds of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

For intranasal administration or administration by inhalation, the active compounds of the invention are conveniently delivered in the form of a solution or suspension from a pump spray container that is squeezed or pumped by the patient or as an aerosol spray presentation from a pressurized container or a nebulizer, with

the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container or nebulizer may contain a solution or suspension of the active compound. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

A proposed dose of the active compounds of the invention for oral, parenteral or buccal administration to the average adult human for the treatment of the conditions referred to above (e.g., rheumatoid arthritis) is 0.1 to 1000 mg of the active ingredient per unit dose which could be administered, for example, 1 to 4 times per day.

Aerosol formulations for treatment of the conditions referred to above (e.g., asthma) in the average adult human are preferably arranged so that each metered dose or "puff" of aerosol contains 20 μ g to 1000 μ g of the compound of the invention. The overall daily dose with an aerosol will be within the range 0.1 mg to 1000 mg. Administration may be several times daily, for example 2, 3, 4 or 8 times, giving for example, 1, 2 or 3 doses each time.

A compound of formula (I) administered in a pharmaceutically acceptable form either alone or in combination with one or more additional agents which modulate a mammalian immune system or with antiinflammatory agents, agents which may include but are not limited to cyclosporin A (e.g. Sandimmune® or Neoral®, rapamycin, FK-506 (tacrolimus), leflunomide, deoxyspergualin, mycophenolate (e.g. Cellcept®, azathioprine (e.g. Imuran®), daclizumab (e.g. Zenapax®), OKT3 (e.g. Orthocolone®), AtGam, aspirin, acetaminophen, ibuprofen, naproxen, piroxicam, and antiinflammatory steroids (e.g. prednisolone or dexamethasone); and such agents may be administered as part of the same or separate dosage forms, via the same or different routes of administration, and on the same or different administration schedules according to standard pharmaceutical practice.

FK506 (Tacrolimus) is given orally at 0.10-0.15 mg/kg body weight, every 12 hours, within first 48 hours postoperative. Does is monitored by serum Tacrolimus trough levels.

Cyclosporin A (Sandimmune oral or intravenous formulation, or Neoral®, oral solution or capsules) is given orally at 5 mg/kg body weight, every 12 hours within 48 hours postoperative. Dose is monitored by blood Cyclosporin A trough levels.

5 The active agents can be formulated for sustained delivery according to methods well known to those of ordinary skill in the art. Examples of such formulations can be found in United States Patents 3,538,214, 4,060,598, 4,173,626, 3,119,742, and 3,492,397.

10 The ability of the compounds of formula I or their pharmaceutically acceptable salts to inhibit Janus Kinase 3 and, consequently, demonstrate their effectiveness for treating disorders or conditions characterized by Janus Kinase 3 is shown by the following *in vitro* assay tests.

Biological Assay

JAK3 (JH1:GST) Enzymatic Assay

15 The JAK3 kinase assay utilizes a protein expressed in baculovirus-infected SF9 cells (a fusion protein of GST and the catalytic domain of human JAK3) purified by affinity chromatography on glutathione-Sepharose. The substrate for the reaction is poly-Glutamic acid-Tyrosine (PGT (4:1), Sigma catalog # P0275), coated onto Nunc Maxi Sorp plates at 100 µg/ml overnight at 37°C. The morning after coating, the plates are washed three times and JAK3 is added to the wells containing
20 100 µl of kinase buffer (50 mM HEPES, pH 7.3, 125 mM NaCl, 24 mM MgCl₂) + 0.2 µM ATP + 1 mM Na orthovanadate.) The reaction proceeds for 30 minutes at room temperature and the plates is washed three more times. The level of phosphorylated tyrosine in a given well is quantitated by standard ELISA assay utilizing an anti-phosphotyrosine antibody (ICN PY20, cat. #69-151-1).

25 Inhibition of Human IL-2 Dependent T-Cell Blast Proliferation

This screen measures the inhibitory effect of compounds on IL-2 dependent T-Cell blast proliferation *in vitro*. Since signaling through the IL-2 receptor requires JAK-3, cell active inhibitors of JAK-3 should inhibit IL-2 dependent T-Cell blast proliferation.

30 The cells for this assay are isolated from fresh human blood. After separation of the mononuclear cells using Accuspin System-Histopaque-1077 (Sigma # A7054), primary human T-Cells are isolated by negative selection using Lympho-Kwik T (One Lambda, Inc., Cat # LK-50T). T-Cells are cultured at 1-2 x 10⁶/ml in Media (RPMI + 10% heat-inactivated fetal calf serum (Hyclone Cat # A-1111-L) + 1%

Penicillin/Streptomycin (Gibco)) and induce to proliferate by the addition of 10ug/ml PHA (Murex Diagnostics, Cat # HA 16). After 3 days at 37°C in 5% CO₂, cells are washed 3 times in Media, resuspended to a density of 1-2 x 10⁶ cells/ml in Media plus 100 Units/ml of human recombinant IL-2 (R&D Systems, Cat # 202-IL). After 1 week
5 the cells are IL-2 dependent and can be maintained for up to 3 weeks by feeding twice weekly with equal volumes of Media + 100 Units/ml of IL-2.

To assay for a test compounds ability to inhibit IL-2 dependent T-Cell proliferation, IL-2 dependent cells are washed 3 times, resuspended in media and then plated (50,000 cells/well/0.1ml) in a Flat-bottom 96-well microtiter plate (Falcon #
10 353075). From a 10 mM stock of test compound in DMSO, serial 2-fold dilutions of compound are added in triplicate wells starting at 10 uM. After one hour, 10 Units/ml of IL-2 is added to each test well. Plates are then incubated at 37°C, 5% CO₂ for 72 hours. Plates are then pulsed with ³H-thymidine (0.5 uCi/well) (NEN Cat # NET-027A), and incubated an additional 18 hours. Culture plates are then harvested with
15 a 96-well plate harvester and the amount of ³H-thymidine incorporated into proliferating cells is determined by counting on a Packard Top Count scintillation counter. Data is analyzed by plotting the % inhibition of proliferation verses the concentration of test compound. An IC₅₀ value (uM) is determined from this plot.

The following Examples illustrate the preparation of the compounds of the
20 present invention but it is not limited to the details thereof. Commercial reagents were utilized without further purification. THF refers to tetrahydrofuran. DMF refers to N,N-dimethylformamide. Low Resolution Mass Spectra (LRMS) were recorded on either a Hewlett Packard 5989®, utilizing chemical ionization (ammonium), or a Fisons (or Micro Mass) Atmospheric Pressure Chemical Ionization (APCI) platform
25 which uses a 50/50 mixture of acetonitrile/water with 0.1% formic acid as the ionizing agent. Room or ambient temperature refers to 20-25°C.

Example 1

Furan-2-yl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-yl}-methanone

30 Method A

1-Benzyl-4-methyl-pyridinium chloride

To a stirred solution of 4-methylpyridine (26 mL/0.268 mol) in 70 mL of acetone was added 31 mL (0.268 mol) of benzylchloride. The resulting mixture was stirred at 50°C for 18 hours. After cooling to room temperature, the reaction

was filtered, washed with acetone and dried under reduced pressure affording 38 g of the title compound. The filtrate was concentrated under reduced pressure producing an additional 5.6 grams of the title compound (74% combined yield). LRMS: 184.

5 **Method B**

1-Benzyl-4-methyl-1,2,3,6-tetrahydro-pyridine

To a stirred solution of the product from Method A (38 grams/0.171 mol) dissolved in 140 mL of 10:1 ethanol/water at 0°C was added 16 grams (0.427 mol) of sodium borohydride portion-wise over 25 minutes. The resulting mixture stirred for 10 18 hours at room temperature, at which time, the reaction was quenched upon addition of 100 mL of water. The reaction mixture was filtered, the filter cake washed with water and ethylacetate, and the combined filtrates concentrated under reduced pressure to remove the organics. The residue was diluted with water (100 mL) and extracted 3 times with 150 mL with ethylacetate. The combined ethylacetate extracts 15 were dried over Na₂SO₄ and concentrated to dryness in vacuo affording 32 grams (100%) of the title compound as a yellow oil. LRMS: 188 (M+1).

Method C

1-Benzyl-4-methyl-piperidin-3-ol

To a solution of the product from Method B (72.45 grams/0.387 mol) 20 dissolved in 240 mL of THF was added 21.4 grams of NaBH₄ and the mixture cooled to 0 °C. A solution of borontrifluoride etherate (109.4 mL dissolved in 200 mL of THF) was then added dropwise over 1.5 hours. Once added, the reaction mixture was brought to room temperature and stirred for 2 hours. The reaction was again cooled to 0 °C and 29.3 mL of water were added dropwise over 15 minutes followed by 25 dropwise addition of 2N sodium hydroxide (97.5 mL) over 20 minutes. The resulting mixture stirred at 0 °C for 40 minutes and was then brought to room temperature. Hydrogen peroxide (30%) (97.5 mL) was added dropwise at a rate so as not to exceed 50 °C in the reaction mixture (approximately 30 minutes). When the addition was complete, the reaction mixture stirred for 10 minutes, then was cooled to 0 °C. 30 Concentrated hydrochloric acid (97.5 mL) was added over 5 minutes, the reaction mixture was reduced to one third its volume in vacuo, and the pH adjusted to 9-10 with 6N sodium hydroxide (aq). The resulting mixture was extracted three times with ether, the combined ether layers dried over MgSO₄ and evaporated to dryness in

vacuo affording 65.32 grams (79%) of the title compound as yellow oil. LRMS: 206.1 (M+1).

Alternative Method: To a solution of the product from Method B (18.7 grams/0.1 mol) in THF (150mL) was added NaBH₄ (6.5 grams/0.170 mol) at room temperature under N₂. The slurry was cooled to 0°C, and BF₃·OEt₂ (15mL, 16.8 grams/0.118 mol) in THF (25mL) was slowly added through an addition funnel. The addition was kept slow enough to keep the temperature of the reaction mixture below 0°C. After the addition; the reaction mixture was stirred at 0°C for 1 hour and room temperature for 1.5 hours. The reaction was re-cooled to 0°C and water (50mL) was added slowly to destroy the excess borane. The reaction was stirred at room temperature for 2 hours, followed by the addition of Oxone® (110 grams/0.343 mol) in water (500mL) at 0°C. The reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction was quenched upon addition of solid NaHSO₃ until all excess oxidant was destroyed (KI/starch test paper). The pH of the reaction mixture was 1-2. The reaction mixture was then extracted 3 times with 50 mL ethyl acetate, the aqueous layer adjusted to pH 12 with 6 N sodium hydroxide and extracted with ethyl acetate (4 times with 100 mL). The organic layer was washed with brine, dried over Na₂SO₄, and concentrated *in vacuo* affording 19.0 grams (92%) of the title compound as an oil. LRMS: 206.1 (M+1).

20 **Method D**

1-Benzyl-4-methyl-piperidin-3-ol -toluene-4-sulfonic acid salt

To a stirred solution of the product from Method C (65.32 grams/0.318 mol) dissolved in 175 mL of acetone and cooled to 0°C was added a solution of *para*-toluenesulfonic acid monohydrate in 350 mL of acetone (dropwise) over 2 hours and the resulting mixture stirred at 0°C for 1.5 hours. The precipitate was filtered and the filter cake washed with 90 mL of diisopropyl ether. The solid product was then dried *in vacuo* affording 58.55 grams (100%) of the title compound as a white solid. LRMS: 378.5 (M+1).

Method E

30 **1-Benzyl-4-methyl-piperidin-3-one**

To a solution of the product from Method D (9.8 grams/0.026 mol) and 31.7 mL of diisopropylethylamine dissolved in 250 mL of dichloromethane and cooled to

0°C was added (dropwise) 12.4 grams of SO₃pyridine complex dissolved in 153 mL of dimethylsulfoxide over a 40 minute period. Once added, the reaction stirred for 1.5 hours at room temperature and was then quenched upon addition of 200 mL of saturated NaHCO₃ (aq). The dichloromethane was removed in vacuo and the
5 remaining aqueous residue extracted four times with diisopropyl ether (150 mL). The combined ether layers were washed four times with water (100 mL), dried over Na₂SO₄ and concentrated to dryness in vacuo affording 3.81 grams (72.97%) of the title compound as yellow oil. LRMS: 204 (M+1).

Method F

10 (1-Benzyl-4-methyl-piperidin-3-yl)-methyl-amine

To a stirred solution of the product from Method E (3.81 grams/0.019 mol) and 38 mL of 2.0 M methylamine in THF was added 2.2 mL of acetic acid and the resulting mixture stirred at room temperature for 1.5 hours. Triacetoxysodiumborohydride (NaB(OAc)₃H) (7.94 grams/0.038 mol) was added as a
15 solid and the new mixture stirred at room temperature for 18 hours. The reaction was quenched with 2 N hydrochloric acid and the pH adjusted to 1. The reaction mixture was washed two times with ether, the aqueous layer then adjusted to pH of 12 with 6 N sodium hydroxide (aq) and extracted three times with dichloromethane. The combined dichloromethane layers were dried over Na₂SO₄, filtered and evaporated to
20 dryness in vacuo affording 3.51 grams (87.75%) of the title compound as dark yellow oil. LRMS: 219.1 (M+1).

Method G

(1-Benzyl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

25 A mixture of 4-chloropyrrolo[2,3-d]pyrimidine (2.4 grams, 15.9 mmol), prepared by the method of Davoll, J. Am. Chem. Soc., (1960), 82, 131, the product from Method F (1.7 grams, 7.95 mmol) and 10 mL of triethylamine were heated in a sealed tube at 100°C for 4 days. After cooling to room temperature and concentration under reduced pressure, the residue was purified by flash
30 chromatography (silica; 3% methanol in dichloromethane) affording 1.0 grams (38%) of the title compound as a colorless oil. LRMS: 336.1 (M+1).

Method H

Methyl-(4-methyl-piperidin-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

To the product from Method G (0.7 grams, 2.19 mmol) dissolved in 15 mL of ethanol was added 0.5 grams of 20% palladium hydroxide on carbon (50% water) (Aldrich) and the resulting mixture agitated (Parr-Shaker) under an atmosphere of hydrogen (50 psi) at room temperature for 2 days. The Celite filtered reaction mixture was concentrated to dryness in vacuo and the residue purified by flash chromatography (silica; 5% methanol in dichloromethane) affording 0.48 grams (90%) of the title compound. LRMS: 246.1 (M+1).

Method I

[1-(4-Methoxy-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

To a stirred solution of 1 mL of pyridine and 9 mL of dichloromethane was added 40 mg (0.163 mmol) of the product from Method H and 20 L of 4-methoxy-benzenesulfonyl chloride and the resulting mixture stirred at room temperature for 18 hours. The reaction was then quenched upon addition of saturated NaHCO₃ (aq), the organic layer was removed and the aqueous layer extracted with dichloromethane. The dichloromethane layer was dried over Na₂SO₄ and concentrated to dryness in vacuo. The residue was purified by PTLC (silica; 10:1 dichloromethane/methanol) affording 22 mg (32%) of the title compound as a light yellow solid. LRMS: 416.5 (M+1).

The title compounds for examples 2-297 were prepared by a method analogous to that described in Example 1.

Example 2

[1-(4-Methoxy-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 416.

Example 3

(1-Benzenesulfonyl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 386.

Example 4

2-(2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-isoindole-1,3-dione

LRMS: 483.

Example 5

Cyclohexanecarboxylic acid (2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide

5 RMS: 463.

Example 6

2-Chloro-N-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-piperidine-1-sulfonyl}-ethyl)-benzamide

LRMS: 492.

10

Example 7

4-Chloro-N-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-piperidine-1-sulfonyl}-ethyl)-benzamide

LRMS: 492.

Example 8

15 Furan-2-carboxylic acid (2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide

LRMS: 447.

Example 9

20 3-Methoxy-N-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-piperidine-1-sulfonyl}-ethyl)-benzamide

LRMS: 487.

Example 10

Isoxazole-5-carboxylic acid (2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide

25 LRMS: 448.

Example 11

2,4-Difluoro-N-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-piperidine-1-sulfonyl}-ethyl)-benzamide

30 LRMS: 493.

Example 12

3-Chloro-N-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-piperidine-1-sulfonyl}-ethyl)-benzamide

LRMS: 492.

Example 13

3-Fluoro-N-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-
piperidine-1-sulfonyl}-ethyl)-benzamid

5 LRMS: 475.

Example 14

2-Fluoro-N-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-
piperidine-1-sulfonyl}-ethyl)-benzamide

LRMS: 475.

10

Example 15

4-Fluoro-N-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-
piperidine-1-sulfonyl}-ethyl)-benzamide

LRMS: 475.

Example 16

15 N-(2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-
1-sulfonyl}-ethyl)-benzamide

LRMS: 457.

Example 17

20 Cyclopropanecarboxylic acid (2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-
d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide

LRMS: 421.

Example 18

Cyclopentanecarboxylic acid (2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-
d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide

25

LRMS: 449.

Example 19

Cyclopentyl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-
piperidin-1-yl}-methanone

30

LRMS: 342.

Example 20

Tetrahydro-furan-2-carboxylic acid (2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-
d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide

LRMS: 451.

Example 21

Tetrahydro-furan-3-carboxylic acid (2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide

5 LRMS: 451.

Example 22

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-yl}-
(tetrahydro-furan-2-yl)-methanone

LRMS: 344.

10

Example 23

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-yl}-
(tetrahydro-furan-3-yl)-methanone

LRMS: 344.

Example 24

15

Cyclohexanecarboxylic acid (3-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-3-oxo-propyl)-amide

LRMS: 427.

Example 25

2-Cyclopropyl-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-
piperidin-1-yl}-ethanone

20

LRMS: 328.

Example 26

2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-1-
carbonyl}-pyrrolidine-1-carboxylic acid tert-butyl ester

25

LRMS: 443.

Example 27

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-yl}-
pyrrolidin-2-yl-methanone

30

LRMS: 343.

Example 28

1-(2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-
1-carbonyl}-pyrrolidin-1-yl)-ethanone hydrochloride

LRMS: 385.

Example 29

Furan-3-yl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-
piperidin-1-yl}-methanone

5 LRMS: 340.

Example 30

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-yl}-
pyridin-2-yl-methanone

LRMS: 351.

10

Example 31

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-yl}-
phenyl-methanone

LRMS: 350.

Example 32

15 1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-yl}-
2-phenyl-ethanone

LRMS: 364.

Example 33

20 2-Cyclopropyl-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-
piperidin-1-yl}-ethanone hydrochloride

LRMS: 364.

Example 34

2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-1-
carbonyl}-pyrrolidine-1-carboxylic acid tert-butyl ester

25

LRMS: 443.

Example 35

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid benzylamide

30

LRMS: 379.

Example 36

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid phenylamide

LRMS: 365.

Example 37

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid tetrahydro-furan-3-yl ester

5 LRMS: 360.

Example 38

1-(4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-1-carbonyl}-piperidin-1-yl)-ethanone

LRMS: 399.

10

Example 39

2-Cyclopentyl-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-ethanone

LRMS: 356.

Example 40

15 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid cyclohexylamide

LRMS: 371.

Example 41

20 Azetidin-3-yl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone trifluoroacetate

LRMS: 443.

Example 42

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-yl}-pyrrolidin-1-yl-methanone

25 LRMS: 343.

Example 43

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid methyl-phenyl-amide

30 LRMS: 379.

Example 44

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-yl}-morpholin-4-yl-methanone

LRMS: 359.

Example 45

Methyl-(4-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

5 LRMS: 323.

Example 46

Methyl-(4-methyl-1-thiazol-2-yl-piperidin-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 329.

10

Example 47

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid pyridin-3-ylamide

LRMS: 366.

Example 48

15 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-fluoro-phenyl)-amide

LRMS: 383.

Example 49

20 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-nitro-phenyl)-amide

LRMS: 410.

Example 50

25 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methoxy-phenyl)-amide

LRMS: 395.

Example 51

4-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-benzoic acid ethyl ester

30

LRMS: 437.

Example 52

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-piperidin-1-yl-methanone

LRMS: 357.

Example 53

Methyl-(4-methyl-5'-nitro-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

5 LRMS: 368.

Example 54

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (3-fluoro-phenyl)-amide

LRMS: 383.

10

Example 55

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (2,4-difluoro-phenyl)-amide

LRMS: 401.

Example 56

15 Methyl-[4-methyl-1-(pyrrolidine-1-sulfonyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 379.

Example 57

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (3-methoxy-phenyl)-amide

20

LRMS: 395.

Example 58

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (3-nitro-phenyl)-amide

25

LRMS: 410.

Example 59

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-pyrrolidine-2-carboxylic acid methyl ester

30

LRMS: 401.

Example 60

Methyl-[4-methyl-1-(5-nitro-thiazol-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 374.

Example 61

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-carboxylic acid methyl ester

5 LRMS: 381.

Example 62

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl}-methanol

LRMS: 353.

10

Example 63

Methyl-[4-methyl-1-(piperidine-1-sulfonyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino

LRMS: 393.

Example 64

15 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (3-cyano-phenyl)-amide

LRMS: 390.

Example 65

20 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (3,4-difluoro-phenyl)-amide

LRMS: 401.

Example 66

Methyl-[4-methyl-1-(morpholine-4-sulfonyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

25 LRMS: 395.

Example 67

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-chloro-phenyl)-amide

30 LRMS: 399.

Example 68

Methyl-[4-methyl-1-(6-methyl-pyridazin-3-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 338.

Example 69

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-cyano-phenyl)-amide

5 LRMS: 390.

Example 70

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid biphenyl-4-ylamide

LRMS: 441.

10

Example 71

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-trifluoromethyl-phenyl)-amide

LRMS: 433.

Example 72

15 Methyl-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-piperidine-1-sulfonyl}-ethyl)-carbamic acid benzyl ester

LRMS: 501.

Example 73

Cyclopropyl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-piperidin-1-yl}-methanone

20

LRMS: 314.

Example 74

Cyclobutyl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-piperidin-1-yl}-methanone

25

LRMS: 328.

Example 75

Tetrahydro-furan-3-carboxylic acid methyl-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide

30

LRMS: 465.

Example 76

Cyclohexanecarboxylic acid methyl-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide

LRMS: 477.

Example 77

(5,7-Dichloro-1H-indol-2-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

5 LRMS: 458.

Example 78

4-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-benzoic acid

LRMS: 409.

10

Example 79

(1-Benzooxazol-2-yl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 363.

Example 80

15 (1H-Indol-2-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 389.

Example 81

20 (5-Fluoro-1H-indol-2-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 407.

Example 82

(5-Methoxy-3-methyl-benzofuran-2-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

25

LRMS: 434.

Example 83

(5-Chloro-benzofuran-2-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

30

LRMS: 424.

Example 84

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-
(5-nitro-benzofuran-2-yl)-methanone

LRMS: 435.

Example 85

(5-Chloro-2,3-dihydro-benzofuran-2-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

5 LRMS: 426.

Example 86

(4-Hydroxy-piperidin-1-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 373.

10

Example 87

1-(2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-benzofuran-5-yl)-ethanone

LRMS: 432.

Example 88

15 1-(3-Methyl-2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-1H-indol-5-yl)-ethanone

LRMS: 445.

Example 89

20 [1-(5-Chloro-benzothiazol-2-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 413.

Example 90

(3-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-3-aza-bicyclo[3.1.0]hex-6-yl)-carbamic acid tert-butyl ester

25 LRMS: 470.

Example 91

3-(4-Chloro-phenoxy)-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-propan-1-one

30 LRMS: 428.

Example 92

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid pyridin-2-ylamide

LRMS: 366.

Example 93

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-1-carbonyl}-piperidine-4-carboxylic acid amide hydrochloride

5 LRMS: 436.

Example 94

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (5-chloro-pyridin-2-yl)-amide

LRMS: 400.

10

Example 95

3-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-1-carbonyl}-cyclopentanone

LRMS: 356.

Example 96

15 (3-Hydroxy-cyclopentyl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 358.

Example 97

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-1-carbonyl}-cyclohexanone

20

LRMS: 370.

Example 98

3-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-1-carbonyl}-cyclohexanone

25

LRMS: 370.

Example 99

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (5-nitro-pyridin-2-yl)-amide

30

LRMS: 413.

Example 100

[4-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidine-1-carbonyl}-amino)-phenyl]-acetic acid

LRMS: 423.

Example 101

(4-Amino-piperidin-1-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-
amino]-piperidin-1-yl}-methanone hydrochloride

5 LRMS: 408.

Example 102

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (6-methyl-pyridin-2-yl)-amide

LRMS: 380.

10

Example 103

1-Methyl-4-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-
piperidine-1-carbonyl}-pyrrolidin-2-one

LRMS: 371.

Example 104

15

1-Benzyl-3-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)- amino]-
piperidine-1-carbonyl}-pyrrolidin-2-one

LRMS: 447.

Example 105

20 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (5-trifluoromethyl-pyridin-2-yl)-amide

LRMS: 434.

Example 106

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-
cyclohexanecarboxylic acid (4-cyano-phenyl)-amide

25

LRMS: 389.

Example 107

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (4-carbamoyl-phenyl)-amide

30

LRMS: 408.

Example 108

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (4-sulfamoyl-phenyl)-amide

LRMS: 444.

Example 109

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (5-methyl-thiazol-2-yl)-amide

5 LRMS: 386.

Example 110

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (5,6-dichloro-benzothiazol-2-yl)-amide

LRMS: 491.

10

Example 111

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methyl-thiazol-2-yl)-amide

LRMS: 386.

Example 112

15 Azetidin-1-yl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone hydrochloride

LRMS: 365.

Example 113

20 [2-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-thiazol-4-yl]-acetic acid ethyl ester

LRMS: 458.

Example 114

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4,5-dimethyl-thiazol-2-yl)-amide

25 LRMS: 400.

Example 115

[2-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-thiazol-4-yl]-acetic acid

30 LRMS: 430.

Example 116

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid benzothiazol-2-ylamide

LRMS: 422.

Example 117

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid thiazol-2-ylamide

5 LRMS: 372.

Example 118

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [6-(2-dimethylamino-ethylamino)-pyridin-3-yl]-amide

LRMS: 452.

10

Example 119

N-(4-Chloro-phenyl)-2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-acetamide

LRMS: 413.

Example 120

15 N,N-Dimethyl-2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-acetamide

LRMS: 331.

Example 121

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [6-(2-pyrrolidin-1-yl-ethylamino)-pyridin-3-yl]-amide

20

LRMS: 478.

Example 122

{2-[5-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-pyridin-2-yloxy]-ethyl}-carbamic acid tert-butyl ester

25

LRMS: 525.

Example 123

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [6-(2-amino-ethoxy)-pyridin-3-yl]-amide

30

LRMS: 425.

Example 124

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methylsulfamoyl-phenyl)-amide

LRMS: 458.

Example 125

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methanesulfonyl-phenyl)-amide

5

LRMS: 443.

Example 126

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (5-methyl-[1,3,4]thiadiazol-2-yl)-amide

LRMS: 387.

10

Example 127

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methylsulfamoyl-phenyl)-amide hydrochloride

LRMS: 495.

Example 128

15

Methyl-[4-methyl-1-(1-phenyl-ethyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 350.

Example 129

20

(3-Hydroxy-pyrrolidin-1-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 359.

Example 130

25

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid tert-butyl ester

LRMS: 346.

Example 131

30

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [4-(2-dimethylamino-ethyl)-thiazol-2-yl]-amide

LRMS: 443.

Example 132

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid 4-methanesulfonyl-benzylamide

LRMS: 457.

Example 133

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-acetylsulfamoyl-phenyl)-amide

5 LRMS: 486.

Example 134

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-phenyl-ethane-1,2-dione

LRMS: 378.

10

Example 135

Methyl-[4-methyl-1-(6-methylamino-pyrimidin-4-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 353.

Example 136

15 Methyl-[4-methyl-1-(6-pyrrolidin-1-yl-pyrimidin-4-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 393.

Example 137

20 [1-(6-Benzylamino-pyrimidin-4-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 429.

Example 138

N,N-Dimethyl-N'-(6-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-pyrimidin-4-yl)-ethane-1,2-diamine

25 LRMS: 410.

Example 139

[1-(6-Chloro-pyrimidin-4-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

30 LRMS: 358.

Example 140

[1-(2-Fluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 354

Example 141

[1-(2-Chloro-pyrimidin-4-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

5 LRMS: 359.

Example 142

[1-(4-Chloro-pyrimidin-2-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 359.

10

Example 143

Methyl-[4-methyl-1-(2-methylamino-pyrimidin-4-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 353.

Example 144

15 Methyl-[4-methyl-1-(4-pyrrolidin-1-yl-pyrimidin-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 353.

Example 145

20 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (3-methyl-isoxazol-5-yl)-amide

LRMS: 370.

Example 146

25 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (3-methyl-isoxazol-4-yl)-amide

LRMS: 370.

Example 147

30 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (5-methyl-isoxazol-3-yl)-amide

LRMS: 370.

Example 148

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (5-tert-butyl-isoxazol-3-yl)-amide

LRMS: 412.

Example 149

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid isoxazol-3-ylamide

5 LRMS: 356.

Example 150

N-Methyl-3-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-propionamide

LRMS: 331.

10

Example 151

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-propan-2-one

LRMS: 302.

Example 152

15 {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-oxo-acetic acid methyl ester

LRMS: 332.

Example 153

(1-Cyclohexylmethyl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

20

LRMS: 342.

Example 154

[1-(5-Amino-thiazol-2-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

25

LRMS: 344.

Example 155

Methyl-(4-methyl-piperidin-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 246.

30

Example 156

3-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-3-oxo-propionic acid methyl ester

LRMS: 346.

Example 157

(1-Benzenesulfonylmethyl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 400.

5

Example 158

(3-Hydroxy-pyrrolidin-1-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 359.

Example 159

10 1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-propane-1,2-dione

LRMS: 316.

Example 160

15 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (6-sulfamoyl-pyridin-3-yl)-amide

LRMS: 445.

Example 161

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (6-acetylamino-pyridin-3-yl)-amide

20

LRMS: 423.

Example 162

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [4-(2-dimethylamino-ethylsulfamoyl)-phenyl]-amide

LRMS: 515.

25

Example 163

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (6-cyano-pyridin-3-yl)-amide

30

LRMS: 391.

Example 164

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-sulfonic acid pyridin-2-ylamide

LRMS: 479.

Example 165

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [6-(pyrrolidine-1-carbonyl)-pyridin-3-yl]-amide

5 LRMS: 463.

Example 166

2-Imidazol-1-yl-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-ethanone

LRMS: 354.

10

Example 167

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-carboxylic acid methylamide

LRMS: 380.

Example 168

15 {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl}-morpholin-4-yl-methanone

LRMS: 436.

Example 169

20 5-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-pyridine-2-carboxylic acid propylamide

LRMS: 451.

Example 170

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-carboxylic acid amide

25 LRMS: 366.

Example 171

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-carbonitrile

30 LRMS: 348.

Example 172

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [4-(pyrrolidine-1-sulfonyl)-phenyl]-amide

LRMS: 498.

Example 173

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [4-(morpholine-4-sulfonyl)-phenyl]-amide

5 LRMS: 514.

Example 174

(3-Hydroxy-pyrrolidin-1-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 359.

10

Example 175

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [6-(morpholine-4-carbonyl)-pyridin-3-yl]-amide

LRMS: 479.

Example 176

15 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid [6-(morpholine-4-carbonyl)-pyridin-3-yl]-amide

LRMS: 479.

Example 177

20 2-Imidazol-1-yl-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-ethanone

. LRMS: 354.

Example 178

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid isoxazol-3-ylamide

25

LRMS: 356.

Example 179

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (2,5-dimethyl-2H-pyrazol-3-yl)-amide

30

LRMS: 383.

Example 180

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (5-cyclopropyl-2-methyl-2H-pyrazol-3-yl)-amide

LRMS: 409.

Example 181

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (3-methyl-isothiazol-5-yl)-amide

5 LRMS: 386.

Example 182

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-ylmethyl}-benzoic acid

. LRMS: 380.

10

Example 183

Methyl-[4-methyl-5'-(pyrrolidine-1-sulfonyl)-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 456.

Example 184

15 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-sulfonic acid methylamide

LRMS: 416.

Example 185

20 4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-ylmethyl}-benzenesulfonamide

LRMS: 415.

Example 186

N-tert-Butyl-4-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-ylmethyl}-benzenesulfonamide

25 LRMS: 472.

Example 187

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-pyrazol-1-yl-ethanone

30 LRMS: 354.

Example 188

Methyl-[4-methyl-1-(5-nitro-benzooxazol-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 408.

Example 189

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-sulfonic acid (2-hydroxy-ethyl)-amide

5 LRMS: 446.

Example 190

N-tert-Butyl-4-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-ylmethyl}-benzenesulfonamide

LRMS: 471.

10

Example 191

N-Methyl-2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-oxo-acetamide

. LRMS: 331.

Example 192

15 [1-(5-Ethanesulfonyl-benzooxazol-2-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 455.

Example 193

20 Methyl-[4-methyl-1-(5-methyl-benzooxazol-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 377.

Example 194

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (6-chloro-pyridin-3-yl)-amide

25 LRMS: 400.

Example 195

Methyl-(4-methyl-1-quinolin-2-yl-piperidin-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

30 LRMS: 373.

Example 196

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-sulfonic acid amide

LRMS: 402.

Example 197

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-
2-pyrrolidin-1-yl-ethane-1,2-dione

5 LRMS: 371.

Example 198

Methyl-[4-methyl-1-(4-methyl-benzooxazol-2-yl)-piperidin-3-yl]-(7H-
pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 377.

10 Example 199

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-
2-morpholin-4-yl-ethane-1,2-dione

LRMS:387.

Example 200

15 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (6-methanesulfonyl-pyridin-3-yl)-amide

LRMS: 444.

Example 201

20 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (6-methanesulfonyl-pyridin-3-yl)-amide

LRMS: 444.

Example 202

Methyl-[4-methyl-1-(6-nitro-benzooxazol-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-
d]pyrimidin-4-yl)-amine

25 LRMS: 408.

Example 203

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (6-methanesulfonyl-pyridin-3-yl)-amide

30 LRMS: 444.

Example 204

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
carboxylic acid (6-methanesulfonyl-pyridin-3-yl)-amide

LRMS: 444.

Example 205

Methyl-[4-methyl-1-(6-nitro-benzooxazol-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

5 LRMS: 408.

Example 206

Methyl-[4-methyl-1-(toluene-3-sulfonyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 400.

10 Example 207

Methyl-[4-methyl-1-(4-trifluoromethyl-benzenesulfonyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 454.

Example 208

15 (1-Benzothiazol-2-yl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 379.

Example 209

20 [1-(5,7-Dimethyl-benzooxazol-2-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 391.

Example 210

25 2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-benzooxazole-6-carboxylic acid methyl ester

LRMS: 421.

Example 211

Methyl-[4-methyl-1-(6-methyl-benzooxazol-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

30 LRMS: 377.

Example 212

[1-(6-Methoxy-benzooxazol-2-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 393.

Example 213

Methyl-[4-methyl-1-(5-trifluoromethyl-benzothiazol-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

5 LRMS: 447.

Example 214

[1-(5,7-Dichloro-benzooxazol-2-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 432.

10

Example 215

[1-(6-Chloro-pyridine-3-sulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 422.

Example 216

15 [1-(4-Chloro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 421.

Example 217

20 [1-(4-Fluoro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 404.

Example 218

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-benzonitrile

25 LRMS: 411.

Example 219

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-benzenesulfonyl fluoride

30 LRMS: 468.

Example 220

2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-benzonitrile

LRMS: 411.

Example 221

**1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-
2-tetrazol-1-yl-ethanone**

5 LRMS: 356.

Example 222

**Methyl-[4-methyl-1-(2,2,2-trifluoro-ethanesulfonyl)-piperidin-3-yl]-(7H-
pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

LRMS: 392.

10

Example 223

**[1-(2,6-Difluoro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-
pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

LRMS: 422.

Example 224

15 **[1-(4-tert-Butyl-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-
pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

LRMS: 442.

Example 225

20 **[1-(2,4-Difluoro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-
pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

LRMS: 422.

Example 226

**Methyl-[4-methyl-1-(2-trifluoromethyl-benzenesulfonyl)-piperidin-3-yl]-(7H-
pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

25 LRMS: 454.

Example 227

**[1-(3,5-Bis-trifluoromethyl-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-
(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

30 LRMS: 522.

Example 228

**[1-(3,5-Dichloro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-
pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

LRMS: 455.

Example 229

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-benzoic acid

5 LRMS: 431.

Example 230

[1-(6-Chloro-pyridine-3-sulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 422.

10

Example 231

[1-(4-Chloro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 421.

Example 232

15 [1-(4-Fluoro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 404.

Example 233

20 4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-benzonitrile

LRMS: 411.

Example 234

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-benzenesulfonyl fluoride

25 LRMS: 468.

Example 235

2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-benzonitrile

30 LRMS: 411.

Example 236

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-tetrazol-1-yl-ethanone

LRMS: 356.

Example 237

Methyl-[4-methyl-1-(2,2,2-trifluoro-ethanesulfonyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

5 LRMS: 392.

Example 238

[1-(2,6-Difluoro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 422.

10

Example 239

[1-(4-tert-Butyl-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 442.

Example 240

15 [1-(2,4-Difluoro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 422.

Example 241

20 Methyl-[4-methyl-1-(2-trifluoromethyl-benzenesulfonyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 454.

Example 242

[1-(3,5-Bis-trifluoromethyl-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

25 LRMS: 522.

Example 243

[1-(3,5-Dichloro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

30 LRMS: 455.

Example 244

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-benzoic acid

LRMS: 431.

Example 245

(3-Fluoro-phenyl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-
amino]-piperidin-1-yl}-methanone

5 LRMS: 368.

Example 246

Isothiazol-4-yl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-
piperidin-1-yl}-methanone

LRMS: 357.

10

Example 247

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-
thiophen-3-yl-methanone

LRMS: 356.

Example 248

15 {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-(5-
methyl-1H-pyrazol-3-yl)-methanone

LRMS: 354.

Example 249

20 (5-Methyl-isoxazol-3-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-
amino]-piperidin-1-yl}-methanone.

LRMS: 355.

Example 250

25 {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-(5-
methyl-thiophen-2-yl)-methanone

LRMS: 371.

Example 251

(4-Fluoro-phenyl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-
amino]-piperidin-1-yl}-methanone

30 LRMS: 368.

Example 252

Methyl-[4-methyl-1-(3-nitro-benzenesulfonyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-
d]pyrimidin-4-yl)-amine

LRMS: 431.

Example 253

[1-(3-Fluoro-benzenesulfonyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

5 LRMS: 404.

Example 254

(2-Fluoro-phenyl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 368.

10

Example 255

(1,5-Dimethyl-1H-pyrazol-3-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 368.

Example 256

15 {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-(2-methyl-thiazol-4-yl)-methanone

LRMS: 371.

Example 257

20 {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-thiazol-4-yl-methanone

LRMS: 357.

Example 258

25 (4-Methyl-isothiazol-5-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 371.

Example 259

30 2,2-Dimethyl-5-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-oxo-ethyl)-[1,3]dioxolan-4-one

LRMS: 403.

Example 260

2-Cyclopropyl-N-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-acetamide

LRMS: 436.

Example 261

N-(2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-methanesulfonamide

5 LRMS: 432.

Example 262

(3-Hydroxy-pyrrolidin-1-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 359.

10

Example 263

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-ylmethyl}-benzonitrile

LRMS: 362.

Example 264

15 3-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-benzenesulfonyl fluoride

LRMS: 469.

Example 265

20 2,2-Dimethyl-5-(2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-oxo-ethyl)-[1,3]dioxolan-4-one

LRMS: 402.

Example 266

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid benzyl ester

25 LRMS: 381.

Example 267

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-ylmethyl}-benzenesulfonamide

30 LRMS: 416.

Example 268

[1-(1H-Imidazol-2-ylmethyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 326.

Example 269

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid 2-chloro-benzyl ester

5

LRMS: 415.

Example 270

Methyl-[4-methyl-1-(1-methyl-1H-imidazol-2-ylmethyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 340.

10

Example 271

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-phenoxy-ethanone

LRMS: 380.

Example 272

15

2-(4-Fluoro-phenoxy)-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-ethanone

LRMS: 381.

Example 273

20

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid 2,2,2-trichloro-ethyl ester

LRMS: 420.

Example 274

2-(2-Chloro-phenoxy)-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-ethanone

25

LRMS: 415.

Example 275

2-(3-Chloro-phenoxy)-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-ethanone

30

LRMS: 415.

Example 276

2-Methanesulfonyl-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-ethanone

LRMS: 367.

Example 277

2-(1,1-Dioxo-tetrahydro-1H-thiophen-3-yl)-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-ethanone

5 LRMS: 407.

Example 278

Methyl-[4-methyl-1-(1-phenyl-ethyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 351.

10

Example 279

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-(toluene-4-sulfonyl)-ethanone

LRMS: 443.

Example 280

15 2-Hydroxy-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-ethanone

LRMS: 304.

Example 281

20 1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-3-nitro-propan-1-one

LRMS: 347.

Example 282

25 5-(2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-oxo-ethyl)-thiazolidine-2,4-dione

LRMS: 404.

Example 283

3-Hydroxy-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-propan-1-one

30

LRMS: 318.

Example 284

N-(4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-4-oxo-butyl)-methanesulfonamide

LRMS: 410.

Example 285

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid 2,2-dimethyl-propyl ester

5 LRMS: 360.

Example 286

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-(thiazolidine-3-sulfonyl)-ethanone

LRMS: 440.

10

Example 287

(3,4-Dihydroxy-pyrrolidin-1-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone

LRMS: 376.

Example 288

15 4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-thiazolidin-2-one

LRMS: 376

Example 289

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid prop-2-ynyl ester

20

LRMS: 328.

Example 290

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (2-cyano-ethyl)-amide

25

LRMS: 342.

Example 291

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (2-cyano-ethyl)-amide

. LRMS: 342.

30

Example 292

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-cyclohexyl}-ethanone oxime

LRMS: 302.

Example 293

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid cyanomethyl-methyl-amide

LRMS: 342.

5

Example 294

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid isopropyl ester

LRMS: 332.

Example 295

10

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (2-cyano-ethyl)-methyl-amide

LRMS: 356.

Example 296

15

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-ylmethyl}-pyridin-1-ol

LRMS: 355.

Example 297

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-acetonitrile

20

LRMS: 285.

Example 298

[1-(2-Fluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

25

Method J

To a solution of the product from Method H (50 mg, mmols?) dissolved in 5 mL of methanol was added 154 ul (mmols?) of 2-fluoro-benzaldehyde. The resulting mixture stirred at room temperature for 4 hours, at which time, x mg (y mmol) of sodium cyanoborohydride were added and the new mixture stirred at room temperature for 18 hours. The reaction was quenched upon addition of 2 drops of 1N NaOH (aq) and the mixture concentrated under reduced pressure to remove the methanol. The residue was dissolved in chloroform and washed with water. The

30

aqueous layer was back washed three times with chloroform, the combined chloroform extracts dried over MgSO_4 and concentrated to dryness in vacuo. The crude product was then purified by flash chromatography (silica; 2.5% methanol in chloroform) affording 36 mg (47.5%) of the title compound as a white solid. LRMS: 372.4 (M+1).

The title compounds for examples 299-324 were prepared by the method analogous to that described in Example 298.

Example 299

(1-Benzyl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 336.

Example 300

(1-Furan-2-ylmethyl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 326.

Example 301

[1-(4-Methoxy-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 366.

Example 302

[1-(4-Fluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 354.

Example 303

Methyl-(4-methyl-1-pyridin-3-ylmethyl-piperidin-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 337.

Example 304

Methyl-(4-methyl-1-thiazol-2-ylmethyl-piperidin-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 343.

Example 305

Methyl-(4-methyl-1-pyridin-2-ylmethyl-piperidin-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 337.

5

Example 306

Methyl-[4-methyl-1-(1-phenyl-ethyl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 350.

Example 307

10 **(1-Benzyl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

LRMS: 336.

Example 308

15 **(1-Benzyl-4-methyl-piperidin-3-yl)-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

LRMS: 336.

Example 309

20 **3-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-ylmethyl}-benzonitrile**

LRMS: 361.

Example 310

25 **[1-(3-Fluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

LRMS: 354.

Example 311

30 **[1-(3-Methoxy-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine**

LRMS: 366.

Example 312

3-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]- piperidin-1-ylmethyl}-benzoic acid

LRMS: 380.

Example 313

[1-(2-Fluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 354.

5

Example 314

[1-(2,6-Difluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 372.

Example 315

10 Methyl-(4-methyl-1-phenethyl-piperidin-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 350.

Example 316

15 [1-(2,3-Difluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 372.

Example 317

[1-(3,4-Difluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

20

LRMS: 372.

Example 318

[1-(4-Methanesulfonyl-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 414.

25

Example 319

Methyl-{4-methyl-1-[4-(piperidine-1-sulfonyl)-benzyl]-piperidin-3-yl}-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 483.

30

Example 320

[1-(3,5-Difluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 372.

Example 321

[1-(3-Chloro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 371.

5

Example 322

[1-(3,5-Difluoro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 372.

Example 323

10

[1-(3-Chloro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 371.

Example 324


15

[1-(3,5-Dichloro-benzyl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

LRMS: 405.


Claims

1. A compound of the formula



- 5 or the pharmaceutically acceptable salt thereof; wherein

R¹ is a group of the formula




wherein y is 0, 1 or 2;

- R⁴ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₁-C₆)alkylsulfonyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl wherein the alkyl, alkenyl and alkynyl groups are optionally substituted by deuterium, hydroxy, amino, trifluoromethyl, (C₁-C₄)alkoxy, (C₁-C₆)acyloxy, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, cyano, nitro, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl or (C₁-C₆)acylamino; or R⁴ is (C₃-C₁₀)cycloalkyl wherein the cycloalkyl group is optionally substituted by deuterium, hydroxy, amino, trifluoromethyl, (C₁-C₆)acyloxy, (C₁-C₆)acylamino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, cyano, cyano(C₁-C₆)alkyl, trifluoromethyl(C₁-C₆)alkyl, nitro, nitro(C₁-C₆)alkyl or (C₁-C₆)acylamino;

- R⁵ is (C₂-C₉)heterocycloalkyl wherein the heterocycloalkyl groups must be substituted by one to five groups consisting of carboxy, cyano, amino, deuterium, hydroxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo, (C₁-C₆)acyl, (C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH, (C₁-C₆)alkylamino-CO-, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkylamino, amino(C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl, (C₁-C₆)acyloxy(C₁-C₆)alkyl, nitro, cyano(C₁-C₆)alkyl, halo(C₁-C₆)alkyl, nitro(C₁-C₆)alkyl, trifluoromethyl, trifluoromethyl(C₁-C₆)alkyl, (C₁-C₆)acylamino, (C₁-C₆)acylamino(C₁-C₆)alkyl, (C₁-C₆)alkoxy(C₁-C₆)acylamino, amino(C₁-C₆)acyl, amino(C₁-C₆)acyl(C₁-C₆)alkyl, (C₁-C₆)alkylamino(C₁-C₆)acyl, ((C₁-C₆)alkyl)₂amino(C₁-C₆)acyl, R¹⁵R¹⁶N-CO-O-, R¹⁵R¹⁶N-CO-(C₁-C₆)alkyl, (C₁-C₆)alkyl-

$S(O)_m$, $R^{15}R^{16}NS(O)_m$, $R^{15}R^{16}NS(O)_m$ (C₁-C₆)alkyl, $R^{15}S(O)_mR^{16}N$, $R^{15}S(O)_mR^{16}N$ (C₁-C₆)alkyl wherein m is 0, 1 or 2 and R^{15} and R^{16} are each independently selected from hydrogen or (C₁-C₆)alkyl; and a group of the formula II



II

5 wherein a is 0, 1, 2, 3 or 4;

b, c, e, f and g are each independently 0 or 1;

d is 0, 1, 2, or 3;

X is $S(O)_n$ wherein n is 0, 1 or 2; oxygen, carbonyl or $-C(=N\text{-cyano})-$;

Y is $S(O)_n$ wherein n is 0, 1 or 2; or carbonyl; and

10 Z is carbonyl, $C(O)O-$, $C(O)NR-$ wherein R is hydrogen or (C₁-C₆)alkyl; or Z is $S(O)_n$ wherein n is 0, 1 or 2;

R^6 , R^7 , R^8 , R^9 , R^{10} and R^{11} are each independently selected from the group consisting of hydrogen or (C₁-C₆)alkyl optionally substituted by deuterium, hydroxy, amino, trifluoromethyl, (C₁-C₆)acyloxy, (C₁-C₆)acylamino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, cyano, cyano(C₁-C₆)alkyl, trifluoromethyl(C₁-C₆)alkyl, nitro, nitro(C₁-C₆)alkyl or (C₁-C₆)acylamino;

R^{12} is (C₆-C₁₀)aryl, (C₂-C₉)heteroaryl, (C₃-C₁₀)cycloalkyl or (C₂-C₉)heterocycloalkyl, wherein the aryl, heteroaryl, cycloalkyl and heterocycloalkyl groups are optionally substituted by one to four groups consisting of hydrogen, deuterium, amino, halo, oxo, hydroxy, nitro, carboxy, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, trifluoromethyl, trifluoromethoxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₃-C₁₀)cycloalkyl, (C₁-C₆)alkyl-CO-NH-, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkoxy, carboxy, carboxy(C₁-C₆)alkyl, carboxy(C₁-C₆)alkoxy, benzyloxycarbonyl(C₁-C₆)alkoxy, (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkoxy, (C₆-C₁₀)aryl, amino, amino(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonylamino, (C₆-C₁₀)aryl(C₁-C₆)alkoxycarbonylamino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₁-C₆)alkylamino(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino(C₁-C₆)alkyl, hydroxy, (C₁-C₆)alkoxy, carboxy, carboxy(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonyl, (C₁-

C₆)alkoxycarbonyl(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-, cyano,
 (C₅-C₉)heterocycloalkyl, amino-CO-NH-, (C₁-C₆)alkylamino-CO-NH-, ((C₁-
 C₆)alkyl)₂amino-CO-NH-, (C₆-C₁₀)arylamino-CO-NH-, (C₅-C₉)heteroarylamino-CO-
 NH-, (C₁-C₆)alkylamino-CO-NH-(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino-CO-NH-(C₁-
 5 C₆)alkyl, (C₆-C₁₀)arylamino-CO-NH-(C₁-C₆)alkyl, (C₅-C₉)heteroarylamino-CO-NH-(C₁-
 C₆)alkyl, (C₁-C₆)alkylcyano, (C₁-C₆)alkylcarboxy(C₁-C₆)alkoxy, (C₁-C₆)alkylcarboxy,
 sulfonylamino, aminosulfonyl, sulfonylamino(C₁-C₆)alkyl, sulfonylaminocarboxy(C₁-
 C₆)alkyl, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-
 C₆)alkyl, (C₆-C₁₀)arylsulfonyl, (C₆-C₁₀)arylsulfonylamino, (C₆-C₁₀)arylsulfonylamino(C₁-
 10 C₆)alkyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₃-
 C₁₀)cycloalkyl, (C₃-C₁₀)cycloalkoxy, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₆-
 C₁₀)arylamino, (C₁-C₆)alkylthio, (C₆-C₁₀)arylthio, (C₁-C₆)alkylsulfinyl, (C₆-
 C₁₀)arylsulfinyl, (C₁-C₆)alkylsulfonyl, (C₆-C₁₀)arylsulfonyl, (C₁-C₆)acyl, (C₁-C₆)alkoxy-
 CO-NH-, (C₁-C₆)alkylamino-CO-, (C₅-C₉)heteroaryl, (C₂-C₉)heterocycloalkyl or (C₆-
 15 C₁₀)aryl wherein the heteroaryl, heterocycloalkyl and aryl groups which are optionally
 substituted on R¹² may be further substituted by one to three groups consisting of
 halo, (C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-NH-, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-
 (C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkoxy,
 carboxy, carboxy(C₁-C₆)alkyl, carboxy(C₁-C₆)alkoxy, benzyloxycarbonyl(C₁-C₆)alkoxy,
 20 (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkoxy, (C₆-C₁₀)aryl, amino, amino(C₁-C₆)alkyl, (C₁-
 C₆)alkoxycarbonylamino, (C₆-C₁₀)aryl(C₁-C₆)alkoxycarbonylamino, (C₁-C₆)alkylamino,
 ((C₁-C₆)alkyl)₂amino, (C₁-C₆)alkylamino(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino(C₁-C₆)alkyl,
 hydroxy, (C₁-C₆)alkoxy, carboxy, carboxy(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonyl, (C₁-
 C₆)alkoxycarbonyl(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-, cyano,
 25 (C₅-C₉)heterocycloalkyl, amino-CO-NH-, (C₁-C₆)alkylamino-CO-NH-, ((C₁-
 C₆)alkyl)₂amino-CO-NH-, (C₆-C₁₀)arylamino-CO-NH-, (C₅-C₉)heteroarylamino-CO-
 NH-, (C₁-C₆)alkylamino-CO-NH-(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino-CO-NH-(C₁-
 C₆)alkyl, (C₆-C₁₀)arylamino-CO-NH-(C₁-C₆)alkyl, (C₅-C₉)heteroarylamino-CO-NH-(C₁-
 C₆)alkyl, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylsulfonylamino, (C₁-
 30 C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₆-C₁₀)arylsulfonyl, (C₆-C₁₀)arylsulfonylamino, (C₆-
 C₁₀)arylsulfonylamino(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonylamino, (C₁-
 C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₅-C₉)heteroaryl and (C₂-C₉)heterocycloalkyl;

R² and R³ are each independently selected from the group consisting of
 hydrogen, deuterium, amino, halo, hydroxy, nitro, carboxy, (C₂-C₆)alkenyl, (C₂-

C₆)alkynyl, trifluoromethyl, trifluoromethoxy, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₃-C₁₀)cycloalkyl wherein the alkyl, alkoxy or cycloalkyl groups are optionally substituted by one to three groups selected from halo, hydroxy, carboxy, amino (C₁-C₆)alkylthio, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₅-C₉)heteroaryl, (C₂-C₉)heterocycloalkyl, (C₃-C₉)cycloalkyl or (C₆-C₁₀)aryl; or R² and R³ are each independently (C₃-C₁₀)cycloalkyl, (C₃-C₁₀)cycloalkoxy, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₆-C₁₀)arylamino, (C₁-C₆)alkylthio, (C₆-C₁₀)arylthio, (C₁-C₆)alkylsulfinyl, (C₆-C₁₀)arylsulfinyl, (C₁-C₆)alkylsulfonyl, (C₆-C₁₀)arylsulfonyl, (C₁-C₆)acyl, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkylamino-CO-, (C₅-C₉)heteroaryl, (C₂-C₉)heterocycloalkyl or (C₆-C₁₀)aryl wherein the heteroaryl, heterocycloalkyl and aryl groups are optionally substituted by one to three halo, (C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-NH-, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-(C₁-C₆)alkoxy, carboxy, carboxy(C₁-C₆)alkyl, carboxy(C₁-C₆)alkoxy, benzyloxycarbonyl(C₁-C₆)alkoxy, (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkoxy, (C₆-C₁₀)aryl, amino, amino(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonylamino, (C₆-C₁₀)aryl(C₁-C₆)alkoxycarbonylamino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₁-C₆)alkylamino(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino(C₁-C₆)alkyl, hydroxy, (C₁-C₆)alkoxy, carboxy, carboxy(C₁-C₆)alkyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-NH-, (C₁-C₆)alkyl-CO-NH-, cyano, (C₅-C₉)heterocycloalkyl, amino-CO-NH-, (C₁-C₆)alkylamino-CO-NH-, ((C₁-C₆)alkyl)₂amino-CO-NH-, (C₆-C₁₀)arylamino-CO-NH-, (C₅-C₉)heteroarylamino-CO-NH-, (C₁-C₆)alkylamino-CO-NH-(C₁-C₆)alkyl, ((C₁-C₆)alkyl)₂amino-CO-NH-(C₁-C₆)alkyl, (C₆-C₁₀)arylamino-CO-NH-(C₁-C₆)alkyl, (C₅-C₉)heteroarylamino-CO-NH-(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₆-C₁₀)arylsulfonyl, (C₆-C₁₀)arylsulfonylamino, (C₆-C₁₀)arylsulfonylamino(C₁-C₆)alkyl, (C₁-C₆)alkylsulfonylamino, (C₁-C₆)alkylsulfonylamino(C₁-C₆)alkyl, (C₅-C₉)heteroaryl or (C₂-C₉)heterocycloalkyl;

with the proviso that R⁵ must be substituted by the group of formula II.

2. A compound according to claim 1, wherein R⁵ is (C₂-C₉)heterocycloalkyl optionally substituted by one to three groups selected from deuterium, hydroxy, (C₁-C₆)alkyl, halo, (C₁-C₆)alkoxy and a group of formula II.

3. A compound according to claim 1, wherein a is 0; b is 1; X is carbonyl; c is 0; d is 0; e is 0; f is 0; and g is 0.

4. A compound according to claim 1, wherein a is 0; b is 1; X is carbonyl; c is 0; d is 1; e is 0; f is 0, and g is 0.
5. A compound according to claim 1, wherein a is 0; b is 1; X is carbonyl; c is 1; d is 0; e is 0; f is 0; and g is 0.
- 5 6. A compound according to claim 1, wherein a is 0; b is 1; X is C(=N=cyano)- ; c is 1; d is 0; e is 0; f is 0; and g is 0.
7. A compound according to claim 1, wherein a is 0; b is 0; c is 0; d is 0; e is 0; f is 0; g is 1; and Z is -C(O)-O- .
8. A compound according to claim 1, wherein a is 0; b is 1; X is S(O)_n ; n is 2; c is 0; d is 0; e is 0; f is 0; and g is 0.
- 10 9. A compound according to claim 1, wherein a is 0; b is 1; X is S(O)_n ; n is 2; c is 0; d is 2; e is 0; f is 1; g is 1; and Z is carbonyl.
10. A compound according to claim 1, wherein a is 0; b is 1; X is S(O)_n ; n is 2; c is 0; d is 2; e is 0; f is 1; and g is 0.
- 15 11. A compound according to claim 1, wherein a is 0; b is 1; X is carbonyl; c is 1; d is 0; e is 1; Y is S(O)_n ; n is 2; f is 0; and g is 0.
12. A compound according to claim 1, wherein a is 0; b is 1; X is S(O)_n ; n is 2; c is 1; d is 0; e is 0; f is 0; and g is 0.
13. A compound according to claim 1, wherein a is 1; b is 1; X is carbonyl; c is 1; d is 0; e is 0; f is 0; and g is 0.
- 20 14. A compound according to claim 1, wherein a is 0; b is 1; X is S(O)_n ; c is 0; d is 1; e is 1; Y is S(O)_n ; n is 2; f is 0; and g is 0.
15. A compound according to claim 1, wherein a is 0; b is 1; X is S(O)_n ; c is 0; d is 2, 3 or 4; e is 1; Y is S(O)_n ; n is 2; f is 1; and g is 0.
- 25 16. A compound according to claim 1, wherein a is 0; b is 1; X is oxygen; c is 0; d is 2, 3 or 4; e is 1; Y is S(O)_n ; n is 2; f is 1; and g is 0.
17. A compound according to claim 1, wherein a is 0; b is 1; X is oxygen; c is 0; d is 2, 3 or 4; e is 1; Y is S(O)_n ; n is 2; f is 0; and g is 0.
18. A compound according to claim 1, wherein a is 0; b is 1; X is carbonyl; c is 1; d is 2, 3 or 4; e is 1; Y is S(O)_n ; f is 0; and g is 0.
- 30 19. A compound according to claim 1, wherein a is 0; b is 1; X is carbonyl; c is 1; d is 2, 3 or 4; e is 1; Y is S(O)_n ; n is 2; f is 1; and g is 0.
20. A compound according to claim 1, wherein R^{12} is $(\text{C}_6\text{-C}_{10})\text{aryl}$ or $(\text{C}_2\text{-C}_9)\text{heteroaryl}$ wherein the aryl or heteroaryl group is optionally substituted by one to

four groups consisting of hydrogen, halo, hydroxy, carboxy, trifluormethyl, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkyl-CO-NH-, amino, amino(C₁-C₆)alkyl, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, cyano, amino-CO-NH-, (C₁-C₆)alkylamino-CO-NH-, ((C₁-C₆)alkyl)₂amino-CO-NH-, (C₅-C₉)heteroaryl-amino-CO-NH-, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkylsulfonylamino, (C₆-C₁₀)arylsulfonylamino, (C₁-C₆)alkylsulfonylamino, and (C₁-C₆)alkoxy-CO-NH-.

21. A compound according to claim 1, wherein said compound is selected from the group consisting of:

4-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-ylmethyl}-benzenesulfonamide;

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-sulfamoyl-phenyl)-amide;

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-nitro-phenyl)-amide;

1-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-tetrazol-1-yl-ethanone;

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methylsulfamoyl-phenyl)-amide;

(3-Hydroxy-pyrrolidin-1-yl)-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-methanone;

[2-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-thiazol-4-yl]-acetic acid;

Methyl-(4-methyl-5'-nitro-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-3-yl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;

5-(2-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-2-oxo-ethyl)-thiazolidine-2,4-dione;

{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-thiazolidin-3-yl-methanone;

Methyl-[4-methyl-1-(5-nitro-thiazol-2-yl)-piperidin-3-yl]-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;

[2-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-thiazol-4-yl]-acetic acid ethyl ester;

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methanesulfonyl-phenyl)-amide;

- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid thiazol-2-ylamide;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-cyano-phenyl)-amide;
- 5 {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-pyrrolidin-1-yl-methanone;
- Furan-2-carboxylic acid (2-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-sulfonyl}-ethyl)-amide;
- {4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}
- 10 (tetrahydro-furan-3-yl)-methanone;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid isoxazol-3-ylamide;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (6-cyano-pyridin-3-yl)-amide;
- 15 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-carbonitrile
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (4-methyl-thiazol-2-yl)-amide;
- 2-Cyclopropyl-1-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-
- 20 amino]-piperidin-1-yl}-ethanone;
- Cyclopentyl-{4-methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidin-1-yl}-Methanone;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid (3-methyl-isoxazol-4-yl)-amide;
- 25 [4-({4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-amino)-phenyl]-acetic acid;
- [1-(5-Amino-thiazol-2-yl)-4-methyl-piperidin-3-yl]-methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-
- 30 carboxylic acid (3-methyl-isothiazol-5-yl)-amide;
- 3-{4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carbonyl}-cyclopentanone;
- 4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid benzyl-methyl-amide; and

4-Methyl-3-[methyl-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]-piperidine-1-carboxylic acid dimethylamide.

22. A pharmaceutical composition for (a) treating or preventing a disorder or condition selected from organ transplant rejection, xeno transplation, lupus, multiple sclerosis, rheumatoid arthritis, psoriasis, Type I diabetes and complications from diabetes, cancer, asthma, atopic dermatitis, autoimmune thyroid disorders, ulcerative colitis, Crohn's disease, Alzheimer's disease, leukemia and other autoimmune diseases or (b) the inhibition of protein kinases or Janus Kinase 3 (JAK3) in a mammal, including a human, comprising an amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof, effective in such disorders or conditions and a pharmaceutically acceptable carrier.

23. A pharmaceutical composition for (a) treating or preventing a disorder or condition selected from organ transplant rejection, xeno transplation, lupus, multiple sclerosis, rheumatoid arthritis, psoriasis, Type I diabetes and complications from diabetes, cancer, asthma, atopic dermatitis, autoimmune thyroid disorders, ulcerative colitis, Crohn's disease, Alzheimer's disease, leukemia and other autoimmune diseases or (b) the inhibition of protein kinases or Janus Kinase 3 (JAK3) in a mammal, including a human, comprising an amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof, alone or in combination with one or more additional agents which modulate a mammalian immune system or with antiinflammatory agents, effective in such disorders or conditions and a pharmaceutically acceptable carrier.

24. A method for the inhibition of protein kinases or Janus Kinase 3 (JAK3) in a mammal, including a human, comprising administering to said mammal an effective amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof.

25. A method for treating or preventing a disorder or condition selected from organ transplant rejection, xeno transplation, lupus, multiple sclerosis, rheumatoid arthritis, psoriasis, Type I diabetes and complications from diabetes, cancer, asthma, atopic dermatitis, autoimmune thyroid disorders, ulcerative colitis, Crohn's disease, Alzheimer's disease, leukemia and other autoimmune diseases in a mammal, including a human, comprising administering to said mammal an amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof, effective in treating such a condition.

26. A method for the inhibition of protein kinases or Janus Kinase 3 (JAK3) in a mammal, including a human, comprising administering to said mammal an effective amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof alone or in combination with one or more additional agents which modulate a mammalian immune system or with antiinflammatory agents.

27. A method for treating or preventing a disorder or condition selected from organ transplant rejection, xeno transplation, lupus, multiple sclerosis, rheumatoid arthritis, psoriasis, Type I diabetes and complications from diabetes, cancer, asthma, atopic dermatitis, autoimmune thyroid disorders, ulcerative colitis, Crohn's disease, Alzheimer's disease, leukemia and other autoimmune diseases in a mammal, including a human, comprising administering to said mammal an amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof, alone or in combination with one or more additional agents which modulate a mammalian immune system or with anti-inflammatory agents, effective in treating such a condition.

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/IB 01/00975

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/505 A61P17/06 A61P19/02 A61P37/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 65909 A (PFIZER PRODUCTS INC.) 23 December 1999 (1999-12-23) page 9, line 20 -page 10, line 24; claims; examples ---	1-27
Y	WO 99 65908 A (PFIZER PRODUCTS INC.) 23 December 1999 (1999-12-23) page 10, line 39 -page 12, line 7; claims; examples ---	1-27
Y	EP 0 795 556 A (PHARMACIA AND UPJOHN) 17 September 1997 (1997-09-17) page 6, line 40 -page 7, line 15; claims; examples --- -/--	1-27



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

12 October 2001

Date of mailing of the international search report

22/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Helps, I

INTERNATIONAL SEARCH REPORT

Intern: Application No
PCT/IB 01/00975

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 00 17203 A (BASF) 30 March 2000 (2000-03-30) page 18, line 24 -page 19, line 26; claims; examples ---	1-27
E	WO 01 42246 A (PFIZER PRODUCTS INC.) 14 June 2001 (2001-06-14) page 8, line 19 - line 39; claims; examples -----	1-27

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern: I Application No

PCT/IB 01/00975

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9965909	A	23-12-1999	AU 4054599 A BR 9912171 A CN 1305479 T EP 1087971 A1 WO 9965909 A1 NO 20006454 A	05-01-2000 10-04-2001 25-07-2001 04-04-2001 23-12-1999 15-02-2001
WO 9965908	A	23-12-1999	AU 3951899 A BR 9911365 A CN 1305480 T EP 1087970 A1 WO 9965908 A1 NO 20006453 A	05-01-2000 13-03-2001 25-07-2001 04-04-2001 23-12-1999 05-02-2001
EP 795556	A	17-09-1997	EP 0795556 A1 JP 9323995 A	17-09-1997 16-12-1997
WO 0017203	A	30-03-2000	AU 6048499 A EP 1114053 A1 NO 20011356 A WO 0017203 A1	10-04-2000 11-07-2001 16-05-2001 30-03-2000
WO 0142246	A	14-06-2001	AU 1295001 A WO 0142246 A2	18-06-2001 14-06-2001